

CHAPTER 2

CLIMATE AND PHYSICAL GEOGRAPHY OF AUSTRALIA

General description of Australia

This chapter has been prepared by the Bureau of Meteorology, Department of Science and the Environment. It is mainly concerned with the climate of Australia, although some geographic comparisons and a summary of landform features influencing climate have been included together with a summary of atmospheric climate controls.

Position and area

Position. Australia, including Tasmania, comprises a land area of 7,682,300 square kilometres. The land lies between latitudes 10° 41' S. (Cape York) and 43° 39' S. (South Cape, Tasmania) and between longitudes 113° 09' E. (Steep Point) and 153° 39' E. (Cape Byron). The most southerly point on the mainland is South Point (Wilson's Promontory) 39° 08' S. The latitudinal distance between Cape York and South Point, Wilson's Promontory (South East Cape, Tasmania) is about 3,180 kilometres (3,680 kilometres) respectively and the longitudinal distance between Steep Point and Cape Byron is about 4,000 kilometres.

Area of Australia compared with areas of other countries. The area of Australia is almost as great as that of the United States of America (excluding Alaska), about 50 per cent greater than Europe (excluding U.S.S.R.) and 32 times greater than the United Kingdom. The following table shows the area of Australia in relation to areas of other continents and selected countries.

AREAS OF CONTINENTS AND SELECTED COUNTRIES, *circa* 1970
(^{'000} square kilometres)

Country	Area	Country	Area
Continental divisions—		Country—	
Europe (a)	4,936	Australia	7,682
Asia (a)	27,532	Canada	9,976
U.S.S.R. (Europe and Asia)	22,402	Germany, Federal Republic of	248
Africa	30,319	Japan	372
North and Central America and West Indies	24,247	New Guinea (b)	462
South America	17,834	New Zealand	269
Oceania	8,504	United Kingdom	244
		United States of America (c)	9,363
Total, World excluding Arctic and Antarctic continents			
	135,771		

(a) Excludes U.S.S.R., shown below. (b) West Irian is included in other Asia. (c) Includes Hawaii and Alaska.

Land forms

The average altitude of the surface of the Australian land mass is only about 300 metres. Approximately 87 per cent of the total land mass is less than 500 metres and 99.5 per cent is less than 1,000 metres. The highest point is Mount Kosciusko (2,228 metres) and the lowest point is Lake Eyre (~15 metres).

Australia has three major landform features: the western plateau, the interior lowlands and the eastern uplands. The western half of the continent consists of a great plateau of altitude 300 to 600 metres. The interior lowlands include the channel country of southwest Queensland (drainage to Lake Eyre) and the Murray-Darling system to the south. The eastern uplands consist of a broad belt of varied width extending from north Queensland to Tasmania and consisting largely of tablelands, ranges and ridges with only limited mountain areas above 1,000 metres.

The rivers of Australia may be divided into two major classes, those of the coastal plains with moderate rates of fall and those of the central plains with very slight fall. Of the rivers of the northern part of the east coast, the longest are the Burdekin and the Fitzroy in Queensland. The Hunter is the

largest coastal river of New South Wales, and the Murray River, with its great tributary the Darling, drains part of Queensland, the major part of New South Wales, and a large part of Victoria, finally flowing into the arm of the sea known as Lake Alexandrina, on the eastern side of the South Australian coast. The total length of the Murray is about 2,520 kilometres, about 650 being in South Australia and about 1,870 kilometres from South Australia to the source. The Darling from its junction with the Murray to its junction with the Culgoa is 1,390 kilometres. The Upper Darling (1,140 kilometres) incorporates the Barwon which commences at the junction of the Culgoa to its junction with the Weir River and the Macintyre River from its junction with the Weir to its source near Maybole. The rivers of the north-west coast of Australia (Western Australia) e.g. the Murchison, Gascoyne, Ashburton, Fortescue, De Grey, Fitzroy, Drysdale, and Ord are of considerable size. So also are those in the Northern Territory, e.g. the Victoria and Daly, and those on the Queensland side of the Gulf of Carpentaria, such as the Gregory, Leichhardt, Cloncurry, Gilbert, and Mitchell. The rivers of Tasmania have short and rapid courses, as might be expected from the configuration of the country.

The 'lakes' of Australia may be divided into three classes; true permanent lakes; lakes which being very shallow, become mere morasses in dry seasons or even dry up, and finally present a cracked surface of salt and dry mud; and lakes which are really inlets of the ocean, opening into a lake-like expanse. The second class, which are a characteristic of the interior lowlands are of considerable extent. The largest are Lake Eyre 9,500 square kilometres, Lake Torrens 5,900 square kilometres and Lake Gairdner 4,300 square kilometres.

For further information on the landforms and the geographical features of Australia earlier issues of the Year Book should be consulted. The list of special articles, etc., at the end of this volume indicates the nature of the information available and its position in the various issues.

Area, coastline, tropical and temperate zones, and standard times. The areas of the States and Territories and the length of the coastline were determined in 1973, by the Division of National Mapping, Department of National Resources, by manually digitising these features from the 1:250,000 map series of Australia. This means that only features of measurable size at this scale were considered. About 60,000 points were digitised at an approximate spacing of 0.5 kilometres. These points were joined by chords as the basis for calculation of areas and coastline lengths by computer.

The approximate high water mark coastline was digitised and included all bays, ports and estuaries which are open to the sea. In these cases, the shoreline was assumed to be where the seaward boundary of the title of ownership would be. In mangroves, the shoreline was assumed to be on the landward side. Rivers were considered in a similar manner but the decisions were rather more subjective, the line being across the river where it appeared to take its true form.

AREA, COASTLINE, TROPICAL AND TEMPERATE ZONES, AND STANDARD TIMES: AUSTRALIA

NOTE. See paragraphs above for methods of estimating area and coastline.

State or Territory	Estimated area			Percentage of total area in		Standard times	
	Total	Percentage of total area	Length of coastline	Tropical zone	Temperate zone	Meridian selected	Ahead of G.M.T. (a)
	km ²		km				hours
New South Wales	801,600	10.43	1,900	..	100	150°E	(b)10.0
Victoria	227,600	2.96	1,800	..	100	150°E	(b)10.0
Queensland	1,727,200	22.48	7,400	54	46	150°E	10.0
South Australia	984,000	12.81	3,700	..	100	142°30'E	(b)9.5
Western Australia	2,525,500	32.87	12,500	37	63	120°E	8.0
Tasmania	67,800	0.88	3,200	..	100	150°E	(b)10.0
Northern Territory	1,346,200	17.52	6,200	81	19	142°30'E	9.5
Australian Capital Territory	2,400	0.03	35	..	100	150°E	(b)10.0
Australia	7,682,300	100.00	36,735	39	61

(a) Greenwich Mean Time.

(b) Because of 'daylight saving' an hour should be added from late October to early March.

Climate of Australia

General

The following information has been prepared by the Bureau of Meteorology, Department of Science. Previously, this chapter of the Year Book also included information about the physical geography of Australia. The information appeared most recently in Year Book No. 61 of 1975-76.

The climate of Australia is predominantly continental but the insular nature of the land mass is significant in producing modification of the continental pattern.

The island continent of Australia is relatively dry with 50 per cent of the area having a median rainfall of less than 300 millimetres per year and 80 per cent less than 600 millimetres. Extreme minimum temperatures are not as low as those recorded in other continents because of the absence of extensive mountain masses and because of the expanse of ocean to the south. However, extreme maxima are comparatively high, reaching 50°C over the inland, mainly due to the great east-west extent of the continent in the vicinity of the Tropic of Capricorn.

Climatic discomfort, particularly heat discomfort, is significant over most of Australia. During summer, prolonged high temperatures and humidity around the northern coasts and high temperatures over the inland cause physical discomfort. In winter, low temperatures and strong cold winds over the interior and southern areas can be severe for relatively short periods.

Climatic controls

The generally low relief of Australia causes little obstruction to the atmospheric systems which control the climate. A notable exception is the eastern uplands which modify the atmospheric flow.

In the winter half of the year (May–October) anticyclones, or high pressure systems, pass from west to east across the continent and often remain almost stationary over the interior for several days. These anticyclones may extend to 4,000 kilometres along their west-east axes. Northern Australia is then influenced by mild, dry south-east trade winds, and southern Australia experiences cool, moist westerly winds. The westerlies and the frontal systems associated with extensive depressions travelling over the Southern Ocean have a controlling influence on the climate of southern Australia during the winter season, causing rainy periods. Cold outbreaks, particularly in south-east Australia occur when cold air of Southern Ocean origin is directed northwards by intense depressions having diameters up to 2,000 kilometres. Cold fronts associated with the southern depressions, or with secondary depressions over the Tasman Sea, may produce large day-to-day changes in temperature in southern areas, particularly in south-east coastal regions.

In the summer half of the year (November–April) the anticyclones travel from west to east on a more southerly track across the southern fringes of Australia directing easterly winds generally over the continent. Fine, warmer weather predominates in southern Australia with the passage of each anticyclone. Heat waves occur when there is an interruption to the eastward progression of the anticyclone (blocking) and winds back northerly and later north-westerly. Northern Australia comes under the influence of summer disturbances associated with the southward intrusion of warm moist monsoonal air from north of the inter-tropical convergence zone resulting in a hot rainy season.



Tropical cyclones develop over the seas to the north-west and the north-east of Australia in summer between November and April. Their frequency of occurrence and the tracks they follow vary greatly from season to season. On the average, about three Coral Sea cyclones per season directly affect the Queensland coast, and about two Indian Ocean cyclones affect the north-western coast. Tropical cyclones approaching the coast usually produce very heavy rain in coastal areas. Some cyclones move inland, losing intensity but still producing widespread heavy rainfall. Individual cyclonic systems may control the weather over northern Australia for periods extending to three weeks.

Rainfall

Annual. The annual 10, 50 and 90 percentile* rainfall maps are shown on plates 2, 3 and 4 respectively. The area of lowest rainfall is east of Lake Eyre in South Australia, where the median (50 percentile) rainfall is only about 100 millimetres. Murnpeowie, with 70 years of record, has a median annual rainfall of 101 millimetres. Another very low rainfall area is in Western Australia in the Giles-Warburton Range region, which has a median annual rainfall of about 150 millimetres. A vast region extending from the west coast near Shark Bay across the interior of Western Australia and South Australia to south-west Queensland and north-west New South Wales has a median annual rainfall of less than 200 millimetres. This region is not normally exposed to moist air masses for extended periods and rainfall is irregular, averaging only one or two days per month. However, in favourable synoptic situations, which occur infrequently over extensive parts of the region, up to 400 millimetres of rain may fall within a few days and result in widespread flooding.

The region with the highest median annual rainfall is the east coast of Queensland between Cairns and Cardwell, where Tully's median is highest (4,400 millimetres). The mountainous region of western Tasmania also has a high annual rainfall, with 3,600 millimetres at Lake Margaret. In the

* The amounts that are not exceeded by 10, 50 and 90 per cent of all recordings are the 10, 50 and 90 percentiles or the first, fifth and ninth deciles respectively. The 50 percentile is usually called the median.






PLATE 4

mountainous areas of north-east Victoria and some parts of the east coastal slopes there are small pockets with median annual rainfall greater than 2,500 millimetres, but the map scale is too small for these to be shown.

The Snowy Mountains area in New South Wales also has a particularly high rainfall. The highest median annual rainfall isohyet drawn for this region is 3,200 millimetres, and it is likely that small areas have a median annual rainfall approaching 4,000 millimetres on the western slopes above 2,000 metres elevation. (Gaffney 1971 (i)).

The following table shows the area distribution of the median annual rainfall derived from the map in Plate 3.

AREA DISTRIBUTION OF MEDIAN ANNUAL RAINFALL: AUSTRALIA
(Per cent)

Median annual rainfall	W.A.	N.T.	S.A.	Qld	N.S.W.(a)	Vic.	Tas.	Aust.
Under 200 mm . . .	43.5	15.5	74.2	10.2	8.0	29.6
200 to 300 mm . . .	29.6	35.6	13.5	13.0	20.3	6.3	..	22.9
300 .. 400	10.5	9.0	6.8	12.3	19.0	19.2	..	11.2
400 .. 500	4.3	6.6	3.2	13.5	12.4	11.8	..	7.6
500 .. 600	3.1	5.8	1.8	11.6	11.3	14.1	12.2	6.6
600 .. 800	4.6	11.6	0.5	20.5	15.1	24.5	18.2	10.7
800 .. 1,200	3.7	9.6	..	12.6	11.3	17.7	25.0	7.7
Above 1,200	0.7	6.3	..	6.3	2.6	6.4	44.6	3.7
Total	100.0	100.00	100.00	100.00	100.00	100.00	100.00	100.00

(a) Includes Australian Capital Territory.

Seasonal. As outlined under the heading of Climatic Controls, the rainfall pattern is strongly seasonal in character with a winter rainfall regime in the south and a summer regime in the north.

The dominance of rainfall over other climatic elements in determining the growth of specific plants in Australia has led to the development of a climatic classification based on two main parameters. These parameters are median annual rainfall and seasonal rainfall incidence (Gaffney 1971 (ii)). Plate 5, below, is a simplified version of the seasonal rainfall zones arising from this classification.

Evaporation and the concept of rainfall effectiveness are taken into account to some extent in this classification by assigning higher median annual rainfall limits to the summer zones than the corresponding uniform and winter zones. The main features of the seasonal rainfall are:

- (a) marked wet summer and dry winter of northern Australia;
- (b) wet summer and relatively dry winter of south-eastern Queensland and north-eastern New South Wales;
- (c) uniform rainfall in south-eastern Australia—much of New South Wales, parts of eastern Victoria and in southern Tasmania;
- (d) marked wet winter and dry summer of south-west Western Australia and (to a lesser extent) of much of the remainder of southern Australia directly influenced by westerly circulation;
- (e) arid area comprising about half of the continent extending from the north-west coast of Western Australia across the interior and reaching the south coast at the head of the Great Australian Bight.

Variability. The adequate presentation of rainfall variability over an extensive geographical area is difficult. Probably the best measures are found in tables compiled for a number of individual stations in some of the Climatic Survey districts. These tables show the percentage chances of receiving specified amounts of rainfall in monthly, seasonal or annual time spans. Statistical indexes of rainfall variation based on several techniques have been used to compile maps showing main features of the variability of annual rainfall over Australia.




PLATE 5



One index for assessing the variability of annual rainfall is given by the ratio of the 90–10 percentile range to the 50 percentile (median value) i.e. Variability Index = $\left\{ \frac{90 - 10}{50} \right\}$ percentiles.

Variability based on this relationship (Gaffney 1975) is shown in Plate 6, page 16. The region of high to extreme variability shown in Plate 6, lies mostly in the arid zone with summer rainfall incidence, AZ(S), defined on Plate 5, page 14. In the winter rainfall zones the variability is generally low to moderate as exemplified by the south-west of Western Australia. In the tropics, random cyclone visitations cause extremely great variations in rainfall from year to year: at Onslow (Western Australia), annual totals varied from 15 mm in 1912 to 1,085 mm in 1961 and, in the four consecutive years 1921 to 1924, the annual totals were 566, 69, 682 and 55 mm respectively. At Whim Creek (Western Australia), where 747 mm have been recorded in a single day, only 4 mm were received in the whole of 1924. Great variability can also occur in the heavy rainfall areas: at Tully (Queensland), the annual rainfalls have varied from 7,899 mm in 1950 to 2,489 mm in 1961.

Rainday frequency. The average number of days per year with rainfall of 0.2 mm or more is shown in Plate 7, page 16.

The frequency of rain-days exceed 150 per year in Tasmania (with a maximum of over 200 in western Tasmania), southern Victoria, parts of the north Queensland coast and in the extreme south-west of Western Australia. Over most of the continent the frequency is less than 50 rain-days per year. The area of low rainfall with high variability, extending from the north-west coast of Western Australia through the interior of the continent, has less than 25 rain-days per year. In the high rainfall areas of northern Australia the number of rain-days is about 80 per year, but heavier falls occur in this region than in southern regions.

Intensity. The highest rainfall intensities for some localities are shown in the first table on page 17. These figures represent intensities over only small areas around the recording points because turbulence and exposure characteristics of the measuring gauge may vary over a distance of a few metres. The highest 24-hour (9 a.m. to 9 a.m.) falls are listed by States in the second table on page 17. Most of the very high 24-hour falls (above 700 millimetres) have occurred in the coastal strip of Queensland, where a tropical cyclone moving close to mountainous terrain provides ideal conditions for spectacular falls. The highest 24-hour fall (907 millimetres) occurred at Crohamhurst, Queensland, on 3 February 1893.



HIGHEST RAINFALL INTENSITIES IN SPECIFIED PERIODS
(millimetres)

Station	Period of record	Years of complete records	Period in hours				
			1	3	6	12	24
			mm	mm	mm	mm	mm
Adelaide	1897-1974	74	69	133	141	141	141
Alice Springs	1951-1974	22	54	55	74	103	138
Brisbane	1911-1974	61	88	144	182	265	311
Broome	1948-1973	26	72	119	130	172	228
Canberra	1932-1970	35	51	68	71	89	139
Carnarvon	1956-1971	16	32	63	83	95	108
Charleville	1953-1974	22	42	66	75	111	142
Cloncurry	1953-1974	19	59	118	164	173	204
Darwin	1953-1973	18	88	101	109	152	191
Esperance	1963-1973	9	23	45	62	68	79
Hobart	1911-1976	63	28	56	87	117	168
Meekatharra	1953-1973	19	26	67	81	99	112
Melbourne	1878-1976	86	79	83	86	97	130
Mildura	1953-1976	22	49	60	65	65	91
Perth	1946-1974	27	32	38	47	64	93
Sydney	1913-1976	60	97	132	166	190	282
Townsville	1953-1974	20	87	145	165	168	275

Source: Pluviograph records in Bureau of Meteorology archives.

HIGHEST DAILY RAINFALLS
(all years to 1977 inclusive)

State	Station	Date	Amount
			mm
Queensland	Crohamhurst	3.2.1893	907
	Finch Hatton	18.2.1958	878
	Mount Dangar	20.1.1970	869
	Port Douglas	1.4.1911	801
Western Australia	Whim Creek	3.4.1898	747
	Fortescue	3.5.1890	593
New South Wales	Dorrigo	21.2.1954	809
	Cordeaux River	14.2.1898	574
Northern Territory	Roper Valley	15.4.1963	545
	Groote Eylandt	28.3.1953	513
Tasmania	Mathinna	5.4.1929	336
	Cullenswood	22.3.1974	352
Victoria	Balook	18.2.1951	275
	Hazel Park	1.12.1934	267
South Australia	Ardrossan	18.2.1946	206
	Oodnadatta	9.2.1976	200

Thunderstorms and hail. A thunder-day at a given location is a calendar day on which thunder is heard at least once. Plate 8, page 19 shows isopleths (isobronts) of the average annual number of thunder-days which varies from 80 per year near Darwin to less than 10 per year over parts of the southern regions. Convective processes during the summer wet season cause high thunderstorm incidence in northern Australia. The generally high incidence (40-60 annually) over the eastern upland areas is produced mainly by orographic uplift of moist air streams.

Hail, mostly of small size (less than 10 millimetres diameter), occurs with winter/spring cold frontal activity in southern Australia. Summer thunderstorms, particularly over the uplands of eastern Australia, sometimes produce large hail (greater than 10 millimetres diameter). Hail capable of piercing light gauge galvanised iron occurs at irregular intervals and sometimes causes widespread damage.

Snow. Generally, snow covers much of the Australian Alps above 1,500 metres for varying periods from late autumn to early spring. Similarly, in Tasmania the mountains are covered fairly frequently above 1,000 metres in these seasons. The area, depth and duration are highly variable and in the altitude range 500-1,000 metres no snow falls in some years. Snowfalls at levels below 500 metres are occasionally experienced in southern Australia, particularly in the foothill areas of Tasmania and Victoria, but falls are usually light and short-lived. In some seasons parts of the eastern uplands

above 1,000 metres from Victoria to south-eastern Queensland have been covered with snow for several weeks. In ravines around Mt Kosciuszko (2,228 metres) small areas of snow may persist through summer but there are no permanent snowfields.

Temperature

Average temperatures. Average annual air temperatures as shown in Plate 9, page 19 range from 28° C along the Kimberley coast in the extreme north of Western Australia to 4° C in the alpine areas of south-eastern Australia. Although annual temperature may be used for broad comparisons, monthly temperatures are required for detailed analyses.



July is the month with the lowest average temperature in all parts of the continent. The months with the highest average temperature are January or February in the south and December in the north (except in the extreme north and north-west where it is November). The slightly lower temperatures of mid-summer in the north are due to the increase in cloud during the wet season.


Average monthly maxima. Maps of average maximum and minimum temperatures for the months of January and July are shown in Plates 10–13 inclusive, pages 20–21.


In January, average maximum temperatures exceed 35° C over a vast area of the interior and exceed 40° C over appreciable areas of the north-west. The consistently hottest part of Australia is around Marble Bar, Western Australia (150 kilometres south-east of Port Hedland) where the average is 41° C and daily maxima during summer may exceed 40° C consecutively for several weeks at a time.

The marked gradients of isotherms of maximum temperature in summer in coastal areas, particularly along the south and west coasts, are due to the penetration inland of fresh sea breezes initiated by the sharp temperature discontinuities between the land and sea surfaces. There are also gradients of a complex nature in south-east coastal areas caused primarily by the uplands.

In July a more regular latitudinal distribution of average maxima is evident. Maxima range from 30° C near the north coast to 5° C in the alpine areas of the south-east.







PLATES 12 and 13

Extreme maxima. Temperatures have exceeded 45°C at nearly all inland stations more than 150 kilometres from the coast and at many places on the north-west and south coasts. Temperatures have exceeded 50°C at some inland stations and at a few near the coast. It is noteworthy that Eucla on the south coast has recorded 50.7°C, the highest temperature in Western Australia. This is due to the long trajectory over land of hot north-west winds from the Marble Bar area. Although the highest temperature recorded in Australia was 53.1°C at Cloncurry (Queensland), more stations have exceeded 50°C in western New South Wales than in other areas due to the long land trajectory of hot winds from the north-west interior of the continent.

Extreme maximum temperatures recorded at selected stations, including the highest recorded in each State, are shown in the table below.

EXTREME MAXIMUM TEMPERATURES

(All years to 1976 inclusive)

Station	°C	Station	°C
Western Australia—		New South Wales—	
Eucla	50.7	Bourke	52.8
Roebourne	47.8	White Cliffs	51.1
Marble Bar	49.2	Walgett	50.1
Northern Territory—		Wilcannia	50.0
Charlotte Waters (near Finke)	48.2	Menindee	49.7
South Australia—		Australian Capital Territory—	
Oodnadatta	50.7	Canberra	42.2
Kyancutta	49.3	Victoria—	
Queensland—		Mildura	50.8
Cloncurry	53.1	Tasmania—	
Winton	50.7	Bushy Park	40.9
Birdsville	50.0	Hobart	40.8

Extreme minima. The lowest temperatures in Australia have been recorded in the Snowy Mountains, where Charlotte Pass (elevation 1,760 metres) has recorded -22.2°C. Temperatures have fallen below -5°C at most inland places south of the tropics and at some places within a few kilometres of southern coasts. At Eyre, on the south coast of Western Australia, a minimum of -3.9°C has been recorded, and at Swansea, on the east coast of Tasmania, the temperature has fallen as low as -4.4°C.

In the tropics, extreme minima below 0°C have been recorded at many places away from the coasts as far north as Herberton, Queensland (-3.3°C). Even very close to the tropical coastline temperatures have fallen to 0°C, a low recording being -0.8°C for Mackay.

The next table shows extreme minimum temperatures recorded at specified stations, including the lowest recorded in each State.

EXTREME MINIMUM TEMPERATURES

(All years to 1976 inclusive)

Station	°C	Station	°C
Western Australia—		New South Wales—	
Dwellingup	-7.0	Charlotte Pass	-22.2
Booylgoo	-6.7	Kiandra	-20.6
Salmon Gums	-5.4	Kosciusko Hotel	-14.4
Northern Territory—		Cooma	-11.2
Alice Springs	-7.5	Australian Capital Territory—	
Tempe Downs	-6.0	Canberra	-10.0
South Australia—		Victoria—	
Yongala	-8.2	Mount Hotham	-12.8
Kyancutta	-7.0	Omeo	-11.7
Queensland—		Bairnsdale	-7.2
Stanthorpe	-11.0	Tasmania—	
Nanango	-9.3	Oatlands	-12.8
		Bothwell	-12.5

Heat waves. Periods with a number of successive days having a temperature higher than 40°C are relatively common in summer over parts of Australia. With the exception of the north-west coast of Western Australia, however, most coastal areas rarely experience more than three successive days of such conditions. The frequency increases inland, and periods of up to ten successive days have been

recorded at many inland stations. This figure increases in western Queensland and north-western Western Australia to more than twenty days in places. The central part of the Northern Territory and the Marble Bar-Nullagine area of Western Australia have recorded the most prolonged heat waves.

Heat waves are experienced in the coastal areas from time to time. During 11-14 January 1939, for example, a severe heat wave affected south-eastern Australia: Adelaide had a record of 47.6°C on the 12th, Melbourne a record of 45.6°C on the 13th and Sydney a record of 45.3°C on the 14th.

Frost. Frost can cause serious losses in agricultural crops, and numerous climatic studies have been made in Australia relating to specific crops cultivated in local areas. Foley (1945 (i)) made a comprehensive study of the incidence of frost at stations recording minimum temperature. Since Foley's work was published, the number of stations recording minimum temperatures has increased appreciably.

Under calm conditions, overnight temperatures at ground level are often as much as 5°C lower than those measured in the instrument screen (base height 1.1 metre) and differences of 10°C have been recorded. Only a small number of stations measure minima at ground level, the lowest recordings being -15.1°C at Canberra and -14.6°C at Stanthorpe (Queensland). Lower readings may be recorded in alpine areas.

Frost frequency depends on location and orography, and even on minor variations in the contour of the land. The parts of Australia which are most subject to frost are the eastern uplands from north-eastern Victoria to the western Darling Downs in southern Queensland. Most stations in this region experience more than ten nights a month with readings of 0°C (or under) for three to five months of the year. On Tasmania's Central Plateau similar conditions occur for three to six months of the year. Frosts may occur within a few miles of the coasts except on the Northern Territory and most of the north Queensland coasts.




PLATE 14

Regions in which frosts may occur at any time of the year comprise most of Tasmania, large areas of the tablelands of New South Wales, much of inland Victoria, particularly the north-east, and a small part of the extreme south-west of Western Australia. Over most of the interior of the continent,

and on the highlands of Queensland as far north as the Atherton Plateau, frosts commence in April and end in September. Minimum temperatures below 0°C are experienced in most of the subtropical interior in June and July.

The length of the frost period for the year is taken as the number of days between the first and last recording of an air temperature of 2°C or less. The median duration of the frost period in days per year is shown in Plate 14, page 23.

The median frost period over the continent varies from over 200 days per year in the south-eastern uplands areas south of the Hunter Valley to zero in northern Australia. In the southern regions of the continent the annual frost period generally decreases from about 100 days inland to below 50 days towards the coast. However, there are appreciable spatial variations depending mainly on local orography. In Tasmania the frost period exceeds 300 days on the uplands and decreases to 100 days near the coast.

The table below includes the average annual frequency of minima of 2°C or less for a wide selection of stations, particularly those prone to frosts. These data show the high spatial variability of frost frequency across Australia. The south-eastern alpine areas, as represented by Kiandra (elevation 1,400 metres), have a frequency exceeding 200. At Kalgoorlie the average annual frequency is 27, at Alice Springs 33, Charleville 37, Canberra 105 and Melbourne 19.

FROST FREQUENCY

Average annual number of frosty nights (screen minimum $\leq 2^{\circ}\text{C}$) and heavy frosts ($\leq 0^{\circ}\text{C}$)

Station	Period of record	Altitude (metres)	Number of frosty nights	Number of heavy frosts
Adelaide (airport)	1955-70	10	8	1
Alice Springs	1940-71	550	33	11
Ballan (near Ballarat)	1944-64	500	63	20
Birdsville	1957-71	40	7	1
Brisbane (Archerfield airport)	1939-49	10	9	3
Canberra	1939-71	570	105	65
Ceduna	1939-71	20	18	5
Charleville	1942-71	290	37	15
Hobart (Risdon)	1957-70	40	25	5
Kalgoorlie	1939-71	360	27	7
Kiandra	1957-69	1,400	226	176
Loch Valley (E of Melbourne)	1943-59	500	101	53
Melbourne (Essendon airport)	1939-71	80	19	4
Mount Gambier	1942-71	60	33	10
Perth (airport)	1944-71	20	5	0
Walgett	1957-71	130	30	7

The next table shows percentiles (20, 50 and 80) of the annual number of frosts at selected stations. The difference between the 20 and 80 percentile figures relative to the 50 percentile (median) shows that there is great variability in the number of frosts at individual stations from year to year.

FROST VARIABILITY

Annual number of frosty nights ($\leq 2^{\circ}\text{C}$) and heavy frosts ($\leq 0^{\circ}\text{C}$) 20, 50 and 80 percentiles

Station	Period of record	Altitude (metres)	Number of frosty nights			Number of heavy frosts		
			Percentiles			Percentiles		
			20	50	80	20	50	80
Alice Springs	1941-71	550	16	27	37	5	8	14
Bathurst	1957-71	705	83	101	111	51	69	76
Beechworth (SW of Albury)	1957-71	550	51	58	73	16	22	26
Bridgetown	1957-69	155	30	43	53	7	11	19
Canberra	1939-71	570	87	105	116	48	64	75
Charleville	1943-71	290	21	35	45	6	14	19
Dubbo	1957-71	262	39	43	50	10	14	27
Hay	1957-71	93	21	34	37	5	9	13
Kalgoorlie	1942-71	360	15	22	31	2	4	9
Kiandra	1957-68	1,400	206	228	250	163	175	193
Kyancutta	1957-69	58	31	39	40	7	14	20
Mount Gambier	1942-71	60	20	27	34	3	6	13
Mundiwindi	1957-69	575	8	11	29	2	3	11

FROST VARIABILITY—*continued*

Station	Period of record	Altitude (metres)	Number of frosty nights			Number of heavy frosts		
			Percentiles			Percentiles		
			20	50	80	20	50	80
Nhill (near Horsham)	1957-71	129	41	47	58	12	17	26
Oatlands	1957-71	435	85	101	111	38	46	57
Omeo	1957-71	660	115	132	138	59	74	83
Richmond (NW of Sydney)	1953-71	20	23	30	40	6	10	13
Sale	1945-71	5	25	34	45	5	11	17
Swansea	1957-71	8	38	45	61	7	13	19
Wandering (SE of Perth)	1957-69	335	41	57	70	13	25	34
Waratah	1957-71	627	104	117	131	35	44	53
Yongala (E of Port Pirie)	1957-69	515	62	75	90	32	39	52

By convention, a heavy frost is taken as corresponding to a minimum screen temperature of 0°C or less—see the two previous tables. The regions of mainland Australia most prone to heavy frosts are the eastern uplands and adjacent areas extending from Victoria through New South Wales to south-eastern Queensland. Stations above 1,000 metres in altitude in the southern parts of these uplands have more than 100 heavy frosts annually, and in the upland areas below 1,000 metres the annual frequency ranges from 100 to about 20. Over the remainder of southern Queensland, New South Wales and Victoria, although there are great spatial variations, the average annual frequency of heavy frosts typically ranges from about 20 inland to 10 towards the coast. Some places on the coast experience heavy frosts (for example Portland, Victoria, with 3 annually).

In Tasmania, uplands above 1,000 metres have more than 100 heavy frosts annually, and in neighbouring areas the frequency is about 100 decreasing to 20 towards the coasts. Even some coastal stations have a relatively high frequency (Swansea, for example, has 13).

The southern half of Western Australia, the whole of South Australia, and the Alice Springs district of the Northern Territory experience heavy frosts. Differences in annual frequencies between places are great but, in general, the frequency is about 10 inland, decreasing towards the coasts. Some places average more than 20 heavy frosts annually, notably Wandering, Western Australia (21) and Yongala, South Australia (29). At Alice Springs the annual average frequency is 11.

Humidity

For details see Year Book No. 62.

Sunshine, cloud and fog

For details see Year Book No. 62.

Global radiation

Global (short wave) radiation includes that radiation energy reaching the ground directly from the sun and that received indirectly from the sky, scattered downwards by clouds, dust particles, etc.

Plates 15 and 16, page 27, show the average global radiation for the months of January and July. The table below shows the variability of daily global radiation for June and December (1968-72) at selected stations.

GLOBAL RADIATION: VARIABILITY OF DAILY AMOUNTS FOR JUNE AND DECEMBER
(mWh.cm⁻²)
(20, 50 and 80 percentile values in milliwatt hours per square centimetre (1971-1977))

Station	June			December		
	Percentiles			Percentiles		
	20	50	80	20	50	80
Alice Springs	380	440	460	630	820	880
Darwin	500	530	560	440	620	720
Melbourne	110	170	220	490	690	850
Perth	180	250	310	690	850	870
Williamstown	160	260	330	560	780	880

A high correlation exists between daily global radiation (Plates 15 and 16, page 27) and daily hours of sunshine. On the north-west coast around Port Hedland, where average daily global radiation is the highest for Australia (640 milliwatt hours), average daily sunshine is also highest, being approximately 10 hours. Sunshine is more dependent on variations in cloud coverage than is global

radiation, since the latter includes diffuse radiation from the sky as well as direct radiation from the sun. An example is Darwin where in the dry month of July sunshine approaches twice that of the wet (cloudy) month of January but global radiation figures for the two months are comparable.

Evaporation

For details see Year Book No. 62

Winds

The mid-latitude anticyclones are the chief determinants of Australia's two main prevailing wind streams. In relation to the west-east axes of the anticyclones these streams are easterly to the north and westerly to the south. The cycles of development, motion and decay of low pressure systems to the north and south of the anticyclones result in diversity of wind flow patterns. Wind variations are greatest around the coasts where diurnal land and sea breeze effects are important.

Wind roses for the months of January and July at 9 a.m. and 3 p.m. at selected stations are shown in Plates 17-20 inclusive, pages 28-29. The wind roses show the percentage frequency of direction (eight points of compass) and speed ranges of winds.

Orography affects the prevailing wind pattern in various ways such as the channelling of winds through valleys, deflection by mountains and cold air drainage from highland areas. An example of this channelling is the high frequency of north-west winds at Hobart caused by the north-west south-east orientation of the Derwent River Valley.

Average wind speeds and prevailing directions at Australian capitals are included in the climatic tables on pages 32-39. Perth is the windiest capital with an average wind speed of 15.6 kilometres per hour; Canberra is the least windy with an average speed of 5.8 kilometres per hour.

The highest wind speeds and wind gusts recorded in Australia have been associated with tropical cyclones. The highest recorded gust was 246 kilometres per hour during a cyclone at Onslow, Western Australia in 1975 and gusts reaching 200 kilometres per hour have been recorded on several occasions in northern Australia with cyclone visitations. The highest gusts recorded at Australian capitals were 217 kilometres per hour at Darwin and 156 kilometres per hour at Perth.

Estimates of the extreme wind gust expected in a given return period* have been derived for places through Australia (Whittingham, 1964). On this basis, for example, Darwin would have an extreme gust for a return period of 10 years of 140 kilometres per hour, Melbourne 135 and Perth 130.

Floods

Widespread flood rainfall may occur anywhere in Australia but it has a higher incidence in the north and in the eastern coastal areas. It is most economically damaging along the shorter streams flowing from the eastern uplands eastward to the seaboard of Queensland and New South Wales. These flood rains are notably destructive in the more densely populated coastal river valleys of New South Wales—the Tweed, Richmond, Clarence, Macleay, Hunter and Nepean-Hawkesbury—all of which experience relatively frequent flooding. Although chiefly summer rains, they may occur in any season.

The great Fitzroy and Burdekin river basins of Queensland receive flood rains during the summer wet season. Much of the run-off due to heavy rain in north Queensland west of the eastern uplands flows southward through the normally dry channels of the network of rivers draining the interior lowlands into Lake Eyre. This widespread rain may cause floods over an extensive area, but it soon seeps away or evaporates, occasionally reaching the lake in quantity. The Condamine and other northern tributaries of the Darling also carry large volumes of water from flood rains south through western New South Wales to the Murray and flooding occurs along their courses at times.



Flood rains occur at irregular intervals in the Murray-Murrumbidgee system of New South Wales and Victoria, the coastal streams of southern Victoria and the north coast streams of Tasmania.

Droughts



Drought, in general terms, refers to an acute water shortage. This is normally due to rainfall deficiency but with other parameters contributing to the actual water availability. The best single measure of water availability in Australia is rainfall, although parameters such as evaporation and soil moisture are significant, or even dominant, in some situations.

Droughts have severe economic effects in Australia and during the years 1864-1973 inclusive there have been at least eight major droughts affecting the greater part of Australia and at least seven other droughts of lesser severity affecting extensive areas (Foley 1957 (ii)). The droughts of 1895-1903 and 1958-68 were probably the most disastrous in their effects on primary industry.



*Return period is the average period between successive occurrences equal to, or greater than, a given speed. For example the extreme wind gust for a return period of 10 years can be expected to occur once in 10 years on the average.



PLATES 15 and 16



PLATES 17 and 18



PLATES 19 and 20

Gibbs and Maher (1967), having defined a drought year at a certain station as one with the year's rainfall in the first decile range, concluded that the occurrence of areas in the first decile range on annual decile maps for the period 1885-1965 corresponded rather well with drought areas discussed by Foley (1957).

One method of assessing the incidence of rainfall deficiency is the analysis of the distribution of annual rainfalls less than the median (Gaffney 1975). The range between the 50 percentile (median) and the 10 percentile gives a measure of the variation in magnitude of annual rainfalls less than the median. The ratio of this range to the 30 percentile value may be used as an index of rainfall deficiency incidence or drought incidence, i.e.:

$$\text{Index of drought incidence} = \left\{ \frac{50-10}{30} \right\} \text{ percentile}$$

For example, the indexes for Onslow (north-west coast of Western Australia) and similarly, for Cape Otway (south coast of Victoria) are derived thus:

$$\text{Index for Onslow} = \left\{ \frac{222-64}{145} \right\} \text{ mm} = 1.09$$

$$\text{Index for Cape Otway} = \left\{ \frac{865-716}{801} \right\} \text{ mm} = 0.19$$

Plate 21, below, shows the distribution of the index of drought incidence over Australia. The intrusions of high index values from the interior to the central coast of Queensland and across western New South Wales are noteworthy. The extreme values on the north-west coast of Western Australia are among the highest in Australia (e.g. Onslow 1.09) due to the dependence of the rainfall on random cyclone tracks.




PLATE 21

The Bureau of Meteorology commenced the issue of *Drought Reviews* in June 1965. These reviews provide a summary of serious rainfall deficiencies and are issued monthly when serious or severe deficiencies exist in any of the rainfall districts. The deficiency criteria are based on monthly rainfall decile analyses. A review of droughts in Australia to 1968 is included in Year Book No. 54, 1968. Summaries of subsequent drought periods may be obtained from the *Drought Reviews*.

Climatic discomfort

For details *see* Year Book No. 62.

Climatic data for capital cities

The averages for a number of elements determined from long-period observations at the Australian capitals to 1976 inclusive, are given in the following pages. Extremes generally cover all available data to 1976 inclusive, whereas averages may only refer to present sites.

CLIMATIC DATA: PERTH, WESTERN AUSTRALIA

(Lat. 31° 57' S., Long. 115° 51' E. Height above M.S.L. 19.5 metres)

BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 22 metres)						Mean amt evapora- tion (mm)	No. days thun- der	Mean daily amt clouds	
		Aver- age (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction		9 a.m., 3 p.m., 9 p.m. (a)			No. clear days	
					9 a.m.	3 p.m.					
No. of years of record	91	30(b)	76	62	30(b)	30(b)	9	79	30(b)	30(b)	
January	1012.6	17.5	48.2	26/76*	81	E	SSW	280	0.9	2.3	
February	1013.0	17.2	40.8	4/73	113	ENE	SSW	241	0.7	2.5	
March	1015.2	16.2	51.9	28/75	113	E	SSW	214	0.7	2.8	
April	1017.9	13.7	50.7	25/00	101	ENE	SSW	124	0.9	3.4	
May	1017.9	13.5	44.5	8/73	119	NE	WSW	83	1.8	4.3	
June	1017.5	13.5	48.6	17/27	129	N	NW	59	1.8	4.7	
July	1018.8	14.2	53.9	20/26	137	NNE	W	58	1.5	4.5	
August	1018.8	15.1	51.3	15/03	156	N	WNW	75	1.3	4.5	
September	1018.4	15.1	45.9	11/05	109	ENE	SSW	105	0.7	3.9	
October	1017.0	16.1	43.0	6/16	105	SE	SW	158	0.8	3.8	
November	1015.5	17.2	48.2	26/75*	101	E	SW	205	0.8	3.1	
December	1013.4	17.7	44.5	24/75	103	E	SSW	241	0.9	2.6	
Year	Totals	1,843	12.8	..	108
	Averages	1016.3	15.6	E	SSW	3.5	..
	Extremes	53.9	..	156
20/7/26											

(a) Scale 0-8.

(b) Standard thirty years normal (1911-1940).

TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (° Celsius)			Extreme air temperature (° Celsius)		Extreme temperature (° Celsius)		Mean daily hours sun- shine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record	79	79	79	80	80	63(a)	30	78
January	29.5	17.6	23.5	43.7	29/56	80.7	22/14	4.2
February	29.8	17.8	23.7	44.6	8/33	8.7	1/02	78.7
March	27.8	16.5	22.2	41.3	14/22	7.7	8/03	75.0
April	24.4	14.0	19.2	37.6	9/10	4.1	20/14	69.4
May	20.6	11.5	16.1	32.4	2/07	1.3	11/14	63.3
June	18.1	9.9	14.1	28.1	5/75	1.6	22/55	57.5
July	17.3	9.0	13.2	26.3	17/76	1.2	7/16	56.2
August	17.9	9.1	13.5	27.8	21/40	1.9	31/08	62.3
September	19.4	10.1	14.8	32.7	30/18	2.6	6/56	67.5
October	21.2	11.4	16.3	37.3	29/67	4.2	6/68	71.8
November	24.5	13.8	19.2	40.3	24/13	5.6	1/04	75.0
December	27.3	16.1	21.7	42.3	31/68	8.6	29/57	76.0
Year { Averages	23.2	13.1	18.2
Extremes	44.6	1.2	80.7	-3.9	..
				8/2/33	7/7/16	22/1/14	31/5/64	

(a) Records discontinued 1963.

(b) 8/1903 and 16/1967.

(c) 8/1952 and 6/1956.

HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)			Rainfall (millimetres)					Fog mean No. days				
	Rel. hum. (%) at 9 a.m.			Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day					
	Mean	Highest mean	Lowest mean										
No. of years of record	30(a)	30(a)	79	79	100	100	102	100	100	79			
January	14.8	51	63	41	8	3	55	1879	Nil (b)	44	27/79*	0.2	
February	14.7	51	65	43	11	3	166	1955	Nil (b)	87	17/55	0.3	
March	14.7	57	66	46	20	4	145	1934	Nil (b)	77	9/34	0.6	
April	13.4	61	75	51	46	8	149	1926	Nil	1920	67	30/04	0.9
May	12.4	70	81	60	125	14	308	1879	14	1964	76	17/42	1.3
June	11.4	75	85	68	185	17	476	1945	55	1877	99	10/20	1.4
July	10.9	76	88	69	175	18	425	1958	61	1876	76	4/91*	1.6
August	10.7	71	83	62	138	18	318	1945	12	1902	74	14/45	1.0
September	11.6	66	75	58	81	14	199	1923	9	1916	47	18/66	0.4
October	11.7	60	75	52	55	11	200	1890	1	1969	50	4/67	0.4
November	12.7	52	66	41	21	6	71	1916	Nil	1891	39	29/56	0.2
December	13.9	51	63	39	14	4	81	1951	Nil	(b)	47	3/51	0.2
Year {	Totals	879	120	8.1
	Averages	12.7	62
	Extremes	88	39	476	..	Nil	(b)	99	..
							6/1945				10/6/20		

(a) Standard thirty years normal (1911-1940).

(b) Various years.

Figures such as 26/76, 29/56, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk (*) relate to nineteenth century.

CLIMATIC DATA: DARWIN, NORTHERN TERRITORY

(Lat 12° 28' S., Long. 130° 51' E. Height above M.S.L. 30 metres)

BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 36 metres)				Mean amt evaporation (mm)	No. days thunder	Mean daily amt clouds	
		Average (km/h)	Highest mean speed in one day (km/h)	Highest gust speed (km/h)	Prevailing direction 9 a.m. 3 p.m.			9 a.m. 3 p.m.	No. of clear days
No. of years of record	90	20	..	22(b)	..	8	35	35	35
January	1,006.2	9.3	..	106	W NW	225	12.9	5.9	1
February	1,006.3	10.6	..	101	W NW	187	10.2	5.8	1
March	1,007.2	7.5	..	157	W NW	190	10.6	5.2	3
April	1,009.3	8.8	..	67	SE NW	218	4.0	2.9	10
May	1,010.9	9.6	..	62	SE E	223	0.5	2.0	16
June	1,012.2	10.1	..	64	SE E	206	0.0	1.4	19
July	1,012.8	8.9	..	62	SE E	229	0.0	1.3	20
August	1,012.6	8.6	..	72	SE NW	238	0.0	1.1	20
September	1,011.7	8.6	..	64	ENE NW	270	1.0	1.8	16
October	1,010.5	9.8	..	85	NE NW	285	5.3	2.7	9
November	1,008.7	8.6	..	117	NW NW	260	11.8	3.9	4
December	1,006.9	9.8	..	217	NW NW	240	14.2	4.9	2
Year { Totals	2,773	70.5	..	121
Year { Averages	1,009.6	9.2	SE NW	3.2	..
Year { Extremes	217

(a) Scale 0-8.

(b) Several incomplete years.

TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (°Celsius)		Extreme air temperature (°Celsius)			Extreme temperature (°Celsius)		Mean daily hours sunshine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record	90	90	90	92(a)	20.0	26(b)	..	21
January	32.2	25.0	28.6	37.8	2/82*	75.6	26.42	5.9
February	31.9	24.8	28.4	38.3	20/87*	73.2	(c)	5.9
March	32.4	24.8	28.6	38.9	(d)	74.3	23/38	6.8
April	33.1	24.2	28.7	40.0	7/83*	72.8	1/38	8.6
May	32.3	22.4	27.4	39.1	8/84*	71.2	5/20	9.3
June	30.9	20.4	25.7	39.0	17/37*	68.5	2/16	9.7
July	30.4	19.6	25.1	36.7	17/88*	68.9	28/17	9.8
August	31.4	20.8	26.1	37.0	30/71*	69.1	28/16	10.4
September	32.7	23.2	27.9	38.9	20/82*	69.5	(f)	10.0
October	33.6	25.0	29.3	40.5	17/92*	71.4	30/38	9.5
November	33.8	25.3	29.6	39.6	9/84*	77.0	14/37	8.6
December	33.2	25.3	29.3	38.9	20/82*	76.2	26/23	7.1
Year { Averages	32.3	23.3	27.9	8.5
Year { Extremes	40.5	10.4	77.0
				17/10/1892	29/7/1942	14/11/37		

(a) Years 1882-1941 at Post Office, 1942-1966 at Aerodrome; 1967-1971 at Regional office; sites not strictly comparable. (b) Records discontinued 1942. (c) 5/1938 and 23/1938. (d) 26/1883 and 27/1883. (e) Recorded at Darwin Aerodrome. All other Statistics from 1967 to 1971 at Regional Office. (f) 28/1916 and 3/1921.

HUMIDITY, RAINFALL AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)				Fog mean No. days
		Mean	Highest mean	Lowest mean	Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly	
No. of years of record	85(a)	90	57(b)	57(b)	86(c)	74	107(d)	107(d)	35
January	31.1	81	89	69	391	19	746	1906	296
February	31.1	81	88	71	330	18	815	1969	279
March	30.7	80	84	69	260	17	1013	1977	241
April	27.0	72	80	60	103	8	603	1891	158
May	21.8	65	76	49	14	1	299	1968	56
June	18.7	63	75	52	3	0	76	1973	36
July	17.6	62	71	47	1	0	65	1900	43
August	20.6	66	73	53	2	0	84	1947	80
September	24.7	68	73	54	13	2	108	1942	71
October	27.7	68	72	60	50	5	339	1954	95
November	29.3	70	75	62	126	11	399	1938	120
December	30.5	75	83	65	243	16	616	1974	279
Year { Totals	1,536	97
Year { Averages	25.9	71
Year { Extremes	89	47	1013	3/77	296
								7/1/1897	

(a) Records to 1966 at Aerodrome. (b) 1882 to 1938 at Post Office. (c) 1869 to 1962 at Post Office; 8 years missing. (d) Highest or lowest at either Post Office, Aerodrome or Regional Office Sites. (e) Various years. (f) April to October. Various years. Figures such as 2/82, 26/42, etc., indicate in respect of the month of reference, the day and year of occurrence. Dates marked with an asterisk (*) relate to nineteenth century.

CLIMATIC DATA: ADELAIDE, SOUTH AUSTRALIA

(Lat. 34° 46'S., Long. 138° 35'E. Height above M.S.L. 43 metres)

BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 22 metres)						Mean amt evapora- tion (mm)	No. days thun- der	Mean daily amt clouds		No. of clear days
		Aver- age (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction		9 a.m., 3 p.m.			9 a.m., 3 p.m. (a)		
					9 a.m.	3 p.m.						
No. of years of record	119	20(b)	20(b)	60	30(c)	30(c)	9	104	108	61		
January	1,013.2	12.8	32.2	12/70	116	SW	SW	261	1.5	3.0	12.0	
February	1,014.3	12.1	28.8	25/67	106	NE	SW	224	1.1	3.0	10.7	
March	1,017.2	11.4	30.7	24/64	126	S	SW	180	0.8	3.3	10.7	
April	1,019.8	11.4	37.4	10/56	130	NE	SW	126	1.0	4.2	6.8	
May	1,020.1	11.3	37.8	19/53	113	NE	NW	80	1.0	4.7	4.5	
June	1,019.8	11.6	29.7	16/70	108	NE	N	57	0.9	5.0	3.8	
July	1,019.9	11.8	32.9	13/64	148	NE	NW	61	0.8	4.9	3.5	
August	1,019.0	12.8	38.2	8/55	121	NE	SW	76	1.1	4.2	4.7	
September	1,017.6	13.2	34.9	16/65	111	NNE	SW	113	1.3	4.3	5.5	
October	1,016.0	13.6	35.4	1/68	121	NNE	SW	169	1.9	4.2	5.6	
November	1,015.1	13.9	36.3	14/68	130	SW	SW	202	2.0	3.9	6.5	
December	1,013.3	13.5	31.1	18/69	121	SW	SW	247	1.5	3.4	8.8	
Year { Totals	1,795	14.9	..	83.1	
Year { Averages	1,017.1	NE	SW	4.0	..	
Year { Extremes	38.2	8/8/65	148	

(a) Scale 0-8.

(b) Records of cup anemometer.

(c) Standard 30 years normal (1931-1960).

TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (° Celsius)			Extreme air temperature (° Celsius)			Extreme temperature (° Celsius)			Mean daily hours sun- shine		
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass					
No. of years of record	119	119	119	121	121	54(a)	117	94				
January	29.6	16.4	23.0	47.6	12/39	7.3	21/84*	82.3	18/82*	1.8	3/77	9.9
February	29.4	16.6	23.0	45.3	12/99*	7.5	23/18	76.9	10/00	2.1	23/26	9.3
March	26.9	15.1	21.0	43.6	9/34	6.6	21/33	78.9	17/83*	0.1	21/33	7.9
April	22.7	12.7	17.7	37.0	5/38	4.2	15/59*	68.3	1/83*	-3.5	30/77	6.0
May	18.7	10.3	14.5	31.9	4/21	2.7	(b)	64.6	12/79*	-3.6	19/28	4.8
June	15.8	8.3	12.1	25.6	4/57	0.3	(c)	59.3	18/79*	-6.1	24/44	4.2
July	15.0	7.3	11.1	26.6	29/75	0.0	24/08	56.9	26/90*	-5.5	30/29	4.3
August	16.4	7.8	12.1	29.4	31/11	0.2	17/59*	60.0	31/92*	-5.1	11/29	5.3
September	18.9	9.0	14.0	35.1	30/61	0.4	4/58*	71.4	23/82*	-3.9	25/27	6.2
October	22.0	10.9	16.5	39.4	21/22	2.3	20/58*	72.2	30/21	-3.0	22/66	7.2
November	25.2	12.9	19.1	45.3	21/65*	4.9	2/09	74.9	20/78*	-0.6	17/76	8.6
December	27.8	14.9	21.4	45.9	29/31	6.1	(d)	79.8	7/99*	-1.0	19/76*	9.4
Year { Averages	22.4	11.8	17.1	6.9
Year { Extremes	47.6	..	0.0	..	82.3	..	-6.1
				12/1/39	24/7/08	18/1/62	24/6/44					

(a) Discontinued 1934 incomplete 1931-1934.

(b) 26/1895 and 24/04.

(c) 27/1876 and 24/44.

(d) 16/1861 and 4/06.

HUMIDITY, RAINFALL, AND FOG

Month	Vapour pres- sure mean 9 a.m. (mb)	Rainfall (millimetres)								Fog mean no. days			
		Rel. hum. (%) at 9 a.m.			Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day					
		Mean	Highest mean	Lowest mean									
No. of years of record	108	108	108	108	137	137	139	139	139	76			
January	11.9	41	59	29	20	4	84	1941	Nil (a)	58	2/89*	0.0	
February	12.5	44	61	30	21	4	155	1925	Nil (a)	141	7/25	0.0	
March	12.0	47	62	29	24	5	117	1878	Nil (a)	89	5/78*	0.0	
April	11.5	57	72	37	44	9	154	1971	Nil	1945	80	5/60*	0.0
May	10.8	67	77	49	69	13	197	1875	3	1934	70	1/53*	0.4
June	10.0	75	84	63	72	15	218	1916	6	1958	54	1/20	1.1
July	9.5	76	87	66	67	16	138	1890	10	1899	44	10/65*	1.3
August	9.7	70	80	54	62	16	157	1852	8	1944	57	19/51*	0.6
September	10.0	61	72	44	51	13	148	1923	7	1951	40	20/23	0.2
October	10.2	52	67	29	44	11	133	1949	1	1969	57	16/08	0.0
November	10.5	45	64	31	31	8	113	1839	1	1963	75	12/60	0.0
December	11.3	42	56	31	26	6	101	1861	Nil	1904	61	23/13	0.0
Year	Totals	531	120	3.6
	Averages	10.5	56
	Extremes	87	29	218	..	Nil (b)	141
							6/1916				7/2/25		

(a) Various years.

(b) December to April, various years.

Figures such as 3/55, 21/84, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk (*) relate to nineteenth century.

CLIMATIC DATA: SYDNEY, NEW SOUTH WALES

(Lat. 33° 52' S., Long. 151° 12' E. Height above M.S.L. 42 metres)

BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 22 metres)					Mean amt evaporation (mm)	No. days thunder	Mean daily amt clouds	
		Average (km/h)	Highest mean speed in one day (km/h)	Highest gust speed (km/h)	Prevailing direction	9 a.m. 3 p.m.			9 a.m. 3 p.m.	No. clear days
No. of years of record	66	25(b)	25(b)	25(b)	25(b)	25(b)	84(c)	56	114	65
January	1,012.7	12.3	30.3	10/49	150	NE	189	3.3	4.7	4.9
February	1,014.2	11.6	30.3	18/57	101	NE	142	2.5	4.8	4.5
March	1,016.4	10.5	33.3	10/44	93	WNW	142	1.7	4.4	5.7
April	1,018.3	10.2	36.2	24/44	116	W	108	1.3	4.1	7.3
May	1,018.7	10.5	33.8	18/55	101	W	85	0.9	3.9	7.7
June	1,018.9	11.6	36.0	10/47	135	W	58	0.8	4.0	8.0
July	1,018.5	11.5	34.3	20/51	106	W	73	0.8	3.5	10.5
August	1,017.9	12.1	39.6	9/51	109	WNW	112	1.4	3.3	10.4
September	1,017.0	11.6	35.1	23/42	113	WNW	150	1.8	3.5	9.1
October	1,015.1	12.3	39.4	1/57	153	WNW	203	2.7	4.1	6.5
November	1,013.4	12.4	31.9	21/54	114	WNW	190	3.6	4.5	5.2
December	1,012.1	12.3	36.2	11/52	121	NE	225	3.8	4.6	4.8
Year { Totals	1,677	24.7	..	84.7
Year { Averages	1,016.1	11.6	4.2	..
Year { Extremes	39.6	..	153

(a) Scale 0-8. (b) Years 1938-1962 inclusive. (c) Richmond records.

TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (° Celsius)			Extreme air temperature (° Celsius)			Extreme temperature (° Celsius)		Mean daily hours sun- shine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass		
No. of years of record	117	117	117	117	117	84(a)	117	55	
January	25.7	18.3	22.0	45.3	14/39	10.6	18/49	7.2	
February	25.4	18.4	21.9	42.1	8/26	9.6	28/63*	6.8	
March	24.5	17.3	20.9	39.2	3/69*	9.3	14/86*	6.3	
April	22.1	14.5	18.3	33.0	(b)	7.0	27/64*	6.2	
May	19.2	11.2	15.2	30.0	1/19	4.4	30/62*	5.8	
June	16.6	9.1	12.8	26.9	11/31	2.1	22/32	5.2	
July	15.8	7.8	11.8	25.7	22/26	2.2	12/90*	6.2	
August	17.4	8.7	13.1	30.4	24/54	2.7	3/72*	6.8	
September	19.6	10.8	15.2	34.6	26/65	4.9	2/45	7.1	
October	21.9	13.3	17.6	37.4	4/42	5.7	6/27	7.3	
November	23.5	16.3	19.4	40.3	6/46	7.7	1/05	7.6	
December	24.9	17.2	21.1	42.2	20/57	9.1	3/24	7.4	
Year { Averages	21.4	13.6	17.4	6.7	
Year { Extremes	45.3	2.1	76.3	-4.4	..	
				14/1/39	22/6/32	14/2/39	4/7/1893		

(a) Records discontinued 1946. (b) 1/36 and 10/69.

HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)				Fog mean No. days
		Mean	Highest mean	Lowest mean	Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly	
No. of years of record	100	100	100	100	117	117	117	117	55
January	18.8	68	78	58	100	13	388	1911	0.3
February	19.2	70	81	60	115	13	564	1956	0.6
March	18.3	74	85	62	131	14	521	1942	1.5
April	15.0	74	87	63	126	13	622	1861	2.1
May	11.9	75	90	63	123	13	585	1919	3.1
June	10.2	76	89	63	133	12	643	1950	2.7
July	9.6	74	88	59	104	11	336	1950	2.1
August	9.5	68	84	54	81	11	378	1899	1.7
September	11.3	66	79	49	69	11	357	1879	0.9
October	13.0	62	77	46	76	12	283	(a)	0.6
November	15.0	62	79	42	78	12	577	1961	0.5
December	17.6	64	77	51	79	13	402	1920	0.4
Year { Totals	1,215	148	16.3
Year { Averages	14.1	69
Year { Extremes	90	42	643	6/1950	..

(a) 1916 and 1959.

Figures such as 10/49, 28/63, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk (*) relate to nineteenth century.

CLIMATIC DATA: CANBERRA, AUSTRALIAN CAPITAL TERRITORY

(Lat. 35° 19' S., Long. 149° 11' E. Height above M.S.L. 577 metres)

BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 10 metres)						Mean amt evaporation (mm)	No. days thunder	Mean daily amt clouds		No. clear days
		Average (km/h)	Highest mean speed in one day (km/h)	Highest gust speed (km/h)	Prevailing direction		9 a.m.			3 p.m.	9 a.m., 3 p.m., (a)	
No. of years of record	36	44(b)	47(b)	38(c)	36(c)	36(c)	8	36	36	36(d)		
January	1,012.1	6.6	24	24/33	121	NW	242	3.3	4.1	7.7		
February	1,013.1	6.1	25	24/33	104	NW	194	3.0	4.4	6.5		
March	1,016.0	5.3	29	28/42	111	SE	165	1.7	4.2	7.6		
April	1,018.8	5.0	30	8/45	106	NW	109	0.8	4.2	6.9		
May	1,019.0	4.4	21	27/58	104	NW	71	0.4	4.5	6.7		
June	1,021.0	4.8	26	2/30	96	NW	46	0.2	4.6	6.5		
July	1,020.2	5.0	38	7/31	102	NW	54	0.1	4.4	7.0		
August	1,018.5	5.9	25	25/36	113	NW	77	0.8	4.4	6.7		
September	1,017.4	6.0	28	28/34	107	NW	115	1.1	4.1	7.9		
October	1,014.8	6.5	23	12/57	119	NW	165	2.2	4.4	6.1		
November	1,011.9	6.9	28	28/42	128	NW	200	3.3	4.4	5.7		
December	1,010.7	6.9	26	11/38	106	NW	259	3.4	4.1	7.5		
Year { Totals	1,697	20.3	..	82.8		
Year { Averages	1,016.1	5.8	NW	4.3	6.9		
Year { Extremes	38	7/7/31	128		

(a) Scale 0-8. (b) Recorded at Forestry and Timber Bureau, Yarralumla, where a cup anemometer is installed. (c) Recorded at Meteorological office, R.A.A.F. Fairbairn, where a Dines Pressure Tube anemometer is installed. (d) 1940-75. Formerly assessed over 37-year period at Yarralumla.

TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (° Celsius)			Extreme air temperature (° Celsius)			Extreme temperature (° Celsius)		Mean daily hours sun- shine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass		
No. of years of record	36	36	36	38	38	..	26	37	
January	27.5	12.9	20.2	41.4	31/68	1.8	1/56	8.9	
February	26.6	12.6	19.6	42.2	1/68	3.0	16/62	8.2	
March	24.3	10.4	17.3	36.4	9/40	-1.1	24/67	(a) 7.5	
April	19.6	6.5	13.1	32.6	12/68	-3.3	26/72	6.9	
May	14.9	2.8	8.9	24.5	10/67	-7.5	30/76	5.6	
June	12.0	0.8	6.4	20.1	3/57	-8.5	8/57	4.8	
July	11.1	-0.3	5.4	19.7	29/75	-10.0	11/71	5.1	
August	12.6	0.8	6.7	21.7	24/54	-7.8	6/74	6.1	
September	15.8	2.7	9.3	28.6	26/65	-5.6	5/40	7.4	
October	19.0	5.8	12.4	32.7	13/46	-3.3	4/57	7.9	
November	22.2	8.2	15.1	38.8	19/44	-1.8	28/67	8.7	
December	26.0	11.1	18.6	38.8	21/53	1.1	18/64	9.1	
Year { Averages	19.3	6.2	12.7	7.2	
Year { Extremes	42.2	-10.0	..	-15.1	..	
				1/2/68	11/7/71		11/7/71		

(a) 30/58 and 24/67.

HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rel. hum (%) at 9 a.m.			Rainfall (millimetres)				Fog mean No. days					
		Mean	Highest mean	Lowest mean	Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly		Greatest in one day				
No. of years of record	36(a)	36	36	36	36	36	38	38	36	36				
January	13.1	60	75	42	61	8	164	1941	1	1947	95	12/45	1.1	
February	14.0	65	81	53	59	7	148	1977	Nil	1968	69	20/74	1.2	
March	13.1	69	81	53	51	7	312	1950	1	1954	66	5/59	2.8	
April	10.7	75	84	38	50	8	164	1974	2	1946	75	2/59	4.1	
May	8.7	84	96	73	51	9	150	1953	1	1976	96	3/48	7.5	
June	7.1	85	97	73	39	9	126	1956	5	1971	45	25/56	7.6	
July	6.6	84	93	68	38	10	103	1960	4	1970	35	10/57	7.7	
August	7.1	80	92	58	47	12	156	1974	7	1944	48	29/74	5.0	
September	8.1	74	82	55	50	10	116	1970	6	1946	41	16/62	4.1	
October	10.0	67	82	50	73	12	161	1976	2	1977	105	21/59	3.1	
November	10.7	59	76	38	64	10	135	1961	4	1977	64	9/50	1.4	
December	12.3	59	74	43	56	8	215	1947	Nil	1967	87	30/48	0.6	
Year	Totals	639	110	46.2
	Averages	9.3	72
	Extremes	97	38	312	3/50	Nil	(b)	105	21/10/59	..

(a) Formerly assessed over 38-year period at Forestry and Timber Bureau, Yarralumla. (b) 12/67 and 2/68. Data shown in the above tables relate to the Meteorological Office, R.A.A.F., Fairbairn, except where otherwise indicated, and generally cover years up to 1976.

Figures such as 24/33, 31/68, etc., indicate, in respect of the month of reference, the day and year of the occurrence.

CLIMATE AND PHYSICAL GEOGRAPHY OF AUSTRALIA

CLIMATIC DATA: MELBOURNE, VICTORIA

(Lat. 37° 49' S., Long. 144° 58' E. Height above M.S.L. 35 metres)

BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 28 metres)					Mean amt evapora- tion (mm)	No. days thun- der	Mean daily amt clouds	
		Average (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction				9 a.m., 3 p.m., (a)	No. clear days
					9 a.m.	3 p.m.				
No. of years of record	119	36(b)	65	68	57	57	9	68	119	68
January	1,012.8	12.8	34.0	27/41	106	S	S	228	1.7	4.1
February	1,014.3	12.4	30.6	13/47	119	S	S	198	1.9	4.0
March	1,016.8	11.3	29.0	3/61	106	N	S	155	1.3	4.3
April	1,018.9	10.9	33.7	27/71	108	N	S	97	0.7	4.7
May	1,019.1	11.4	33.0	4/61	116	N	N	59	0.4	5.2
June	1,019.0	11.4	36.7	16/47	103	N	N	38	0.2	5.3
July	1,018.6	12.8	36.9	24/70	109	N	N	47	0.2	5.2
August	1,017.5	12.5	34.3	20/42	108	N	N	60	0.6	5.0
September	1,016.0	12.7	34.0	15/64	111	N	S	91	0.8	4.8
October	1,014.7	12.8	30.4	6/68	111	N	S	130	1.6	4.8
November	1,013.9	13.3	35.8	8/71	114	SW	S	161	1.9	4.9
December	1,012.4	13.1	33.8	12/52	100	S	S	209	2.2	4.5
Year { Totals	1,468	13.4	..	48.0
Year { Averages	1,016.2	12.3	N	S	4.7
Year { Extremes	36.9	..	119
			24/7/70							

(a) Scale 0-8.

(b) Early records not comparable.

(c) Records to 1966.

TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (° Celsius)			Extreme air temperature (° Celsius)		Extreme temperature (° Celsius)			Mean daily hours sun- shine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass		
No. of years of record	120	120	120	122	122	86(a)	118	52(b)	
January	25.8	13.9	19.9	45.6	13/39	5.6	28/85*	8.1	
February	25.7	14.2	19.9	43.1	7/01	4.6	24/24	75.3	
March	23.7	12.8	18.3	41.7	11/40	2.8	17/84*	73.6	
April	20.1	10.5	15.3	34.9	5/38	1.6	24/88*	66.7	
May	16.5	8.3	12.4	28.7	7/05	-1.2	29/16	61.4	
June	13.9	6.6	10.3	22.4	2/57	-2.2	11/66	53.9	
July	13.3	5.7	9.5	23.1	30/75	-2.8	21/69*	52.1	
August	14.8	6.4	10.6	25.0	20/85*	-2.1	11/63*	58.6	
September	17.1	7.6	12.4	31.4	28/28	-0.6	3/40	61.2	
October	19.5	9.2	14.4	36.9	24/14	0.1	3/71*	67.9	
November	21.8	10.8	16.3	40.9	27/94*	2.4	2/96*	70.9	
December	24.1	12.6	18.4	43.7	15/76	4.4	4/70*	76.8	
Year { Averages	19.9	9.9	14.8	
Year { Extremes	45.6	-2.8	..	81.4	-6.7	
				13/1/39	21/7/69		14/1/62	30/6/29	

(a) Discontinued 1946.

(b) Discontinued 1967.

(c) 17/1884 and 20/1897.

HUMIDITY, RAINFALL, AND FOG

Month	Vapour pres- sure mean	Rainfall (millimetres)							Fog mean no. days
	9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Mean no. of days of rain	Greatest monthly	Least monthly	Greatest in one day	
		Mean	Highest mean	Lowest mean					
No. of years of record	68	68	70	70	120	120	122	122	118
January	13.1	61	68	50	48	8	176	1963	0.1
February	14.1	63	77	48	50	7	238	1972	0.3
March	13.3	66	79	50	54	9	191	1911	0.7
April	11.7	72	82	66	59	11	195	1960	1.8
May	10.3	79	88	69	57	14	142	1942	3.6
June	9.3	83	92	73	50	14	114	1859	4.6
July	8.9	81	87	73	49	15	178	1891	4.3
August	9.1	75	82	64	50	15	111	1939	2.3
September	9.5	68	76	60	59	14	201	1916	0.8
October	10.5	63	72	52	68	14	193	1869	0.4
November	11.3	61	71	52	59	12	206	1954	0.2
December	12.5	60	69	48	58	10	182	1863	0.2
Year { Totals	661	143	19.3
Year { Averages	11.1	69
Year { Extremes	92	43	238	2/72	..
								Nil	4/23
								108	29/1/63

(a) Less than 1 mm.

Figures such as 27/41, 28/85, etc., indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk (*) relate to nineteenth century.

CLIMATIC DATA: HOBART, TASMANIA

(Lat. 42° 53' S., Long. 147° 20' E. Height above M.S.L. 54 metres)

BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 12 metres)						Mean amt evapora- tion (mm)	No. days thunder	Mean daily amt clouds		
		Aver- age (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction		9 a.m.			3 p.m.	9 a.m., 3 p.m., 9 p.m. (a)	No. of clear days
No. of years of record	90	63	67	92	30(b)	30(b)	10(c)	64	90	30(b)		
January	1,010.6	12.6	33.5	30/16	130	NNW	SSE	167	1.0	5.0		
February	1,012.9	11.5	40.6	4/27	121	NNW	SSE	135	1.0	4.9		
March	1,014.3	11.0	34.4	13/38	127	NW	SSE	109	0.7	4.8		
April	1,015.5	10.9	38.8	9/52	141	NW	W	70	0.3	5.0		
May	1,015.4	10.4	35.4	21/65	135	NNW	NW	38	0.0	5.0		
June	1,015.2	10.2	38.2	27/20	132	NW	NW	22	0.0	5.0		
July	1,014.0	10.7	36.9	22/53	129	NNW	NNW	26	0.0	4.8		
August	1,012.8	10.9	41.0	19/26	140	NNW	NW	44	0.1	4.8		
September	1,011.4	12.5	43.1	28/65	150	NNW	NW	73	0.1	4.9		
October	1,010.3	12.6	32.4	3/65	140	NNW	SW	107	0.4	5.2		
November	1,009.8	12.8	34.1	18/15	135	NNW	S	123	0.6	5.3		
December	1,009.4	12.4	37.7	1/34	122	NNW	SSE	150	0.8	5.3		
Year { Totals	1,064	5.1	..		
Averages	1,012.6	11.5	NNW	W	5.0		
Extremes	43.1	28/9/65	150		

(a) Scale 0-8.

(b) Standard thirty years normal (1911-1940).

(c) Class "A" American pan.

TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (° Celsius)			Extreme air temperature (° Celsius)			Extreme temperature (° Celsius)			Mean daily hours sun- shine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass			
No. of years of record	92	92	92	94	94	57(a)	91	79		
January	21.4	11.5	16.5	40.8	4/76	4.5 (b)	71.1 (c)	-0.8 19/97*	7.9	
February	21.5	11.8	16.7	40.2	12/99*	3.9 20/87*	73.9 24/68*	-2.0 -/87*	7.0	
March	20.0	10.6	15.3	37.3	13/40	1.8 31/26	66.1 26/44	-2.5 30/02	6.4	
April	17.1	8.7	12.9	30.6	1/41	0.6 14/63	61.1 18/93*	-3.9 -/86*	5.0	
May	14.2	6.7	10.5	25.5	5/21	-1.6 30/02	53.3 (d)	-6.7 19/02	4.3	
June	11.8	5.1	8.5	20.6	1/07	-2.8 25/72	50.0 12/94*	-7.7 24/63	3.9	
July	11.4	4.4	7.9	21.0	30/75	-2.4 (f)	49.4 12/93*	-7.4 16/86*	4.3	
August	12.8	5.0	8.9	24.5	26/77	-1.8 5/62	54.4 -/87*	-6.6 7/09	5.0	
September	14.9	6.2	10.6	28.2	29/73	-0.6 16/97*	58.9 23/93*	-7.6 16/26	5.9	
October	16.7	7.5	12.1	33.4	24/14	0.0 12/89*	68.9 9/93*	-4.6 (e)	6.3	
November	18.5	9.0	13.8	36.8	26/37	1.6 16/41	55.6 19/92*	-3.4 1/08	7.0	
December	20.2	10.5	15.4	40.7	30/97*	3.3 3/06	71.9 10/39	-2.6 -/86*	7.2	
Year { Averages	16.7	8.1	12.4	5.9	
Extremes	40.8	-2.8	..	73.9	-7.7	..	
				4/1/1976	25/6/72	..	24/2/1868	24/6/1963	..	

(a) Period 1934-1938 not comparable; records discontinued 1946. (b) 09/1937 and 11/1937. (c) 05/1886 and 13/1905. (d) -/1899 and -/1893. (e) 1/1886 and 1/1899. (f) 11/1895 and 7/1973.

HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rainfall (millimetres)							Fog mean No. days					
		Rel. hum. (%) at 9 a.m.			Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day						
		Mean	Highest mean	Lowest mean										
No. of years of record	77(a)	81	84	84	93	93	95	95	131	64				
January	11.0	58	81	45	49	11	150	1893	4	1958	75	30/16	0.3	
February	11.7	62	83	49	42	10	171	1964	3	1914	56	1/54	0.1	
March	11.0	65	78	52	47	11	255	1946	7	1943	88	17/46	0.3	
April	10.0	70	84	57	55	12	248	1960	2	1904	133	23/60	0.3	
May	8.8	75	86	61	49	14	214	1958	4	1913	47	3/73	1.1	
June	7.9	78	91	61	59	14	238	1954	7	(c)	147	7/54	1.7	
July	7.6	78	87	72	54	15	157	1974	8	1950	64	18/22	1.4	
August	7.9	73	86	59	49	16	161	1946	8	1892	65	2/76	0.8	
September	8.3	66	81	52	52	15	201	1957	10	1931	156	15/57	0.2	
October	9.1	62	74	52	64	17	193	1947	10	1914	66	4/06	0.1	
November	9.6	59	73	49	56	14	188	1885	9	(d)	94	30/85*	0.1	
December	10.6	58	73	42	57	13	196	(b)	5	(e)	85	5/41	0.1	
Year {	Totals	633	162	6.1
	Averages	9.5	67
	Extremes	91	42	255	..	2	..	156
							3/1946		4/1904		15/9/57			

(a) 1894-1970.

(b) 1897 and 1916.

(c) 1886 and 1967.

(d) 1919 and 1921.

(e) 1897, 1915 and 1931.

Figures such as 30/16, 12/99, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk (*) relate to nineteenth century.

REFERENCES

Bureau of Meteorology (Melbourne), Department of Science and the Environment

- Ashton, H. T. 1964 *Meteorological Data for Air Conditioning in Australia*. Bulletin 47.
- Foley, J. C. 1945 (i) *Frost in the Australian Region*. Bulletin 32.
1957 (ii) *Droughts in Australia*. Bulletin 43.
- Gaffney, D. O. 1971 (i) *Snowy Mountains Median Annual Rainfall*. Working Paper 139.
1971 (ii) *Seasonal Rainfall Zones in Australia*. Paper 141.
- Gibbs, W. J. and Maher, J. V. 1967 *Rainfall Deciles as Drought Indicators*. Bulletin 48.
- Hounam, C. E. 1969 *Climate and Air Conditioning Requirements in Sparsely Occupied Areas of Australia*. Working Paper 30/78.
- Whittingham, H. E. . . . 1964 *Extreme Wind Gusts in Australia*. Bulletin 46.

Other Publications

- American Society of Heating, 1960 *Physiological Principles. In Heating, Ventilating and Air Refrigerating and Air Conditioning Guide* Vol. 38 (Washington: the Society).
- Gaffney, D. O. 1970 *Rainfall*.
1973 *Temperatures*
1973 *Climate*
Commentaries prepared for *Atlas of Australian Resources*, Second Series, Department of Minerals and Energy, Canberra.
- 1975 *Rainfall Deficiency and Evaporation in relation to Drought in Australia*. ANZAAS Congress, Canberra.
- Lee, D. H. K. and Henschel, 1963 *Evaluation of Environment in Shelters*. U.S. Department of Health, Education and Welfare (Cincinnati: the Department).