

## CHAPTER 2

# CLIMATE OF AUSTRALIA

### General

The following information has been prepared by the Bureau of Meteorology, Department of Science. Previously, this chapter of the Year Book also included information about the physical geography of Australia. The information appeared most recently in Year Book No. 61 of 1975-76.

The climate of Australia is predominantly continental but the insular nature of the land mass is significant in producing modification of the continental pattern.

The island continent of Australia is relatively dry with 50 per cent of the area having a median rainfall of less than 300 millimetres per year and 80 per cent less than 600 millimetres. Extreme minimum temperatures are not as low as those recorded in other continents because of the absence of extensive mountain masses and because of the expanse of ocean to the south. However, extreme maxima are comparatively high, reaching 50°C over the inland, mainly due to the great east-west extent of the continent in the vicinity of the Tropic of Capricorn.

Climatic discomfort, particularly heat discomfort, is significant over most of Australia. During summer, prolonged high temperatures and humidity around the northern coasts and high temperatures over the inland cause physical discomfort. In winter, low temperatures and strong cold winds over the interior and southern areas can be severe for relatively short periods.

### Climatic controls

The generally low relief of Australia causes little obstruction to the atmospheric systems which control the climate. A notable exception is the eastern uplands which modify the atmospheric flow.

In the winter half of the year (May-October) anticyclones, or high pressure systems, pass from west to east across the continent and often remain almost stationary over the interior for several days. These anticyclones may extend to 4,000 kilometres along their west-east axes. Northern Australia is then influenced by mild, dry south-east trade winds, and southern Australia experiences cool, moist westerly winds. The westerlies and the frontal systems associated with extensive depressions travelling over the Southern Ocean have a controlling influence on the climate of southern Australia during the winter season, causing rainy periods. Cold outbreaks, particularly in south-east Australia occur when cold air of Southern Ocean origin is directed northwards by intense depressions having diameters up to 2,000 kilometres. Cold fronts associated with the southern depressions, or with secondary depressions over the Tasman Sea, may produce large day-to-day changes in temperature in southern areas, particularly in south-east coastal regions.



In the summer half of the year (November-April) the anticyclones travel from west to east on a more southerly track across the southern fringes of Australia directing easterly winds generally over the continent. Fine, warmer weather predominates in southern Australia with the passage of each anticyclone. Heat waves occur when there is an interruption to the eastward progression of the anticyclone (blocking) and winds back northerly and later north-westerly. Northern Australia comes under the influence of summer disturbances associated with the southward intrusion of warm moist monsoonal air from north of the inter-tropical convergence zone, resulting in a hot rainy season.

Tropical cyclones develop over the seas to the north-west and the north-east of Australia in summer between November and April. Their frequency of occurrence and the tracks they follow vary greatly from season to season. On the average, about three Coral Sea cyclones per season directly affect the Queensland coast, and about two Indian Ocean cyclones affect the north-western coast. Tropical cyclones approaching the coast usually produce very heavy rain in coastal areas. Some cyclones move inland, losing intensity but still producing widespread heavy rainfall. Individual cyclonic systems may control the weather over northern Australia for periods extending to three weeks.

### Rainfall

*Annual.* The annual 10, 50 and 90 percentile\* rainfall maps are shown on plates 2, 3 and 4 respectively. The area of lowest rainfall is east of Lake Eyre in South Australia, where the median (50 percentile) rainfall is only about 100 millimetres. Murnpeowie, with 70 years of record, has a median annual rainfall of 101 millimetres. Another very low rainfall area is in Western Australia in the Giles-Warburton Range region, which has a median annual rainfall of about 150 millimetres. A vast region extending from the west coast near Shark Bay across the interior of Western Australia and South Australia to south-west Queensland and north-west New South Wales has a median annual rainfall of less than 200 millimetres. This region is not normally exposed to moist air masses for extended periods and rainfall is irregular, averaging only one or two days per month. However, in favourable synoptic situations, which occur infrequently over extensive parts of the region, up to 400 millimetres of rain may fall within a few days and result in widespread flooding.

\* The amounts that are not exceeded by 10, 50 and 90 per cent of all recordings are the 10, 50 and 90 percentiles or the first, fifth and ninth deciles respectively. The 50 percentile is usually called the median.



PLATES 2 and 3

# CLIMATE OF AUSTRALIA




PLATE 4

The region with the highest median annual rainfall is the east coast of Queensland between Cairns and Cardwell, where Tully's median is highest (4,400 millimetres). The mountainous region of western Tasmania also has a high annual rainfall, with 3,600 millimetres at Lake Margaret. In the mountainous areas of north-east Victoria and some parts of the east coastal slopes there are small pockets with median annual rainfall greater than 2,500 millimetres, but the map scale is too small for these to be shown.

The Snowy Mountains area in New South Wales also has a particularly high rainfall. The highest median annual rainfall isohyet drawn for this region is 3,200 millimetres, and it is likely that small areas have a median annual rainfall approaching 4,000 millimetres on the western slopes above 2,000 metres elevation. (Gaffney 1971 (i)).

The following table shows the area distribution of the median annual rainfall derived from the map in Plate 3.

**AREA DISTRIBUTION OF MEDIAN ANNUAL RAINFALL: AUSTRALIA**  
(Per cent)

Median annual rainfall	W.A.	N.T.	S.A.	Qld	N.S.W.(a)	Vic.	Tas.	Aust.
Under 200 mm .	43.5	15.5	74.2	10.2	8.0	..	..	29.6
200 to 300 mm .	29.6	35.6	13.5	13.0	20.3	6.3	..	22.9
300 " 400 "	10.5	9.0	6.8	12.3	19.0	19.2	..	11.2
400 " 500 "	4.3	6.6	3.2	13.5	12.4	11.8	..	7.6
500 " 600 "	3.1	5.8	1.8	11.6	11.3	14.1	12.2	6.6
600 " 800 "	4.6	11.6	0.5	20.5	15.1	24.5	18.2	10.7
800 " 1,200 "	3.7	9.6	..	12.6	11.3	17.7	25.0	7.7
Above 1,200 "	0.7	6.3	..	6.3	2.6	6.4	44.6	3.7
Total .	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

(a) Includes Australian Capital Territory.

*Seasonal.* As outlined under the heading of Climatic Controls, the rainfall pattern is strongly seasonal in character with a winter rainfall regime in the south and a summer regime in the north.

The dominance of rainfall over other climatic elements in determining the growth of specific plants in Australia has led to the development of a climatic classification based on two main parameters. These parameters are median annual rainfall and seasonal rainfall incidence (Gaffney 1971 (ii)). Plate 5, page 28, is a simplified version of the seasonal rainfall zones arising from this classification.

Evaporation and the concept of rainfall effectiveness are taken into account to some extent in this classification by assigning higher median annual rainfall limits to the summer zones than the corresponding uniform and winter zones. The main features of the seasonal rainfall are:

- (a) marked wet summer and dry winter of northern Australia;
- (b) wet summer and relatively dry winter of south-eastern Queensland and north-eastern New South Wales;
- (c) uniform rainfall in south-eastern Australia—much of New South Wales, parts of eastern Victoria and in southern Tasmania;
- (d) marked wet winter and dry summer of south-west Western Australia and (to a lesser extent) of much of the remainder of southern Australia directly influenced by westerly circulation;
- (e) arid area comprising about half of the continent extending from the north-west coast of Western Australia across the interior and reaching the south coast at the head of the Great Australian Bight.

*Variability.* The adequate presentation of rainfall variability over an extensive geographical area is difficult. Probably the best measures are found in tables compiled for a number of individual stations in some of the Climatic Survey districts. These tables show the percentage chances of receiving specified amounts of rainfall in monthly, seasonal or annual time spans. Statistical indexes of rainfall variation based on several techniques have been used to compile maps showing main features of the variability of annual rainfall over Australia.




PLATE 5



One index for assessing the variability of annual rainfall is given by the ratio of the 90-10 percentile range to the 50 percentile (median value) i.e. Variability Index =  $\left\{ \frac{90 - 10}{50} \right\}$  percentiles.

Variability based on this relationship (Gaffney 1975) is shown in Plate 6, page 30. The region of high to extreme variability shown in Plate 6, lies mostly in the arid zone with summer rainfall incidence, AZ(S), defined in Plate 5, page 28. In the winter rainfall zones the variability is generally low to moderate as exemplified by the south-west of Western Australia. In the tropics, random cyclone visitations cause extremely great variations in rainfall from year to year: at Onslow (Western Australia), annual totals varied from 15 mm in 1912 to 1,085 mm in 1961 and, in the four consecutive years 1921 to 1924, the annual totals were 566, 69, 682 and 55 mm respectively. At Whim Creek (Western Australia), where 747 mm have been recorded in a single day, only 4 mm were received in the whole of 1924. Great variability can also occur in the heavy rainfall areas: at Tully (Queensland), the annual rainfalls have varied from 7,899 mm in 1950 to 2,489 mm in 1961.

*Rainday frequency.* The average number of days per year with rainfall of 0.2 mm or more is shown in Plate 7, page 30.

The frequency of rain-days exceed 150 per year in Tasmania (with a maximum of over 200 in western Tasmania), southern Victoria, parts of the north Queensland coast and in the extreme south-west of Western Australia. Over most of the continent the frequency is less than 50 rain-days per year. The area of low rainfall with high variability, extending from the north-west coast of Western Australia through the interior of the continent, has less than 25 rain-days per year. In the high rainfall areas of northern Australia the number of rain-days is about 80 per year, but heavier falls occur in this region than in southern regions.

*Intensity.* The highest rainfall intensities for some localities are shown in the first table on page 31. These figures represent intensities over only small areas around the recording points because turbulence and exposure characteristics of the measuring gauge may vary over a distance of a few metres. The highest 24-hour (9 a.m. to 9 a.m.) falls are listed by States in the second table on page 31. Most of the very high 24-hour falls (above 700 millimetres) have occurred in the coastal strip of Queensland, where a tropical cyclone moving close to mountainous terrain provides ideal conditions for spectacular falls. The highest 24-hour fall (907 millimetres) occurred at Crohamhurst, Queensland on 3 February 1893.



PLATES 6 and 7

## HIGHEST RAINFALL INTENSITIES IN SPECIFIED PERIODS

(millimetres)

Station	Period of record	Years of complete records	Period in hours				
			1	3	6	12	24
			mm	mm	mm	mm	mm
Adelaide . . .	1897-1967 . . .	67	69	133	141	141	141
Alice Springs . . .	1951-1970 . . .	18	54	55	64	87	106
Brisbane . . .	1911-1968 . . .	14	88	144	182	244	308
Broome . . .	1948-1970 . . .	23	72	119	130	172	228
Canberra . . .	1932-1970 . . .	35	51	68	71	89	138
Carnarvon . . .	1956-1971 . . .	16	32	63	82	95	108
Charleville . . .	1953-1971 . . .	19	42	66	75	111	142
Cloncurry . . .	1953-1972 . . .	17	46	118	164	173	204
Darwin . . .	1953-1970 . . .	15	88	101	109	152	191
Esperance . . .	1963-1972 . . .	8	23	45	62	68	79
Hobart . . .	1911-1976 . . .	63	28	56	87	117	168
Meekatharra . . .	1953-1971 . . .	17	26	67	80	98	112
Melbourne . . .	1878-1976 . . .	86	79	83	86	102	129
Mildura . . .	1953-1976 . . .	22	49	60	65	65	91
Perth . . .	1946-1971 . . .	24	32	38	47	64	93
Sydney . . .	1913-1976 . . .	60	97	139	162	180	281
Townsville . . .	1953-1970 . . .	16	87	111	122	161	275

Source: Pluviograph records in Bureau of Meteorology archives.

## HIGHEST DAILY RAINFALLS

(all years to 1976 inclusive)

State	Station	Date	Amount
			mm
Queensland . . .	Crohamhurst . . .	3.2.1893	907
	Finch Hatton . . .	18.2.1958	878
	Mount Dangar . . .	20.1.1970	869
	Port Douglas . . .	1.4.1911	801
Western Australia . . .	Whim Creek . . .	3.4.1898	747
	Fortescue . . .	3.5.1890	593
New South Wales . . .	Dorrigo . . .	21.2.1954	809
	Cordeaux River . . .	14.2.1898	574
Northern Territory . . .	Roper Valley . . .	15.4.1963	545
	Groote Eylandt . . .	28.3.1953	513
Tasmania . . .	Mathinna . . .	5.4.1929	336
	Cullenswood . . .	22.3.1974	352
Victoria . . .	Balook . . .	18.2.1951	275
	Hazel Park . . .	1.12.1934	267
South Australia . . .	Ardrossan . . .	18.2.1946	206
	Oodnadatta . . .	9.2.1977	200

**Thunderstorms and hail.** A thunder-day at a given location is a calendar day on which thunder is heard at least once. Plate 8, page 33 shows isopleths (isobronts) of the average annual number of thunder-days which varies from 80 per year near Darwin to less than 10 per year over parts of the southern regions. Convective processes during the summer wet season cause high thunderstorm incidence in northern Australia. The generally high incidence (40-60 annually) over the eastern upland areas is produced mainly by orographic uplift of moist air streams.

Hail, mostly of small size (less than 10 millimetres diameter), occurs with winter/spring cold frontal activity in southern Australia. Summer thunderstorms, particularly over the uplands of eastern Australia, sometimes produce large hail (greater than 10 millimetres diameter). Hail capable of piercing light gauge galvanised iron occurs at irregular intervals and sometimes causes widespread damage.

*Snow.* Generally, snow covers much of the Australian Alps above 1,500 metres for varying periods from late autumn to early spring. Similarly, in Tasmania the mountains are covered fairly frequently above 1,000 metres in these seasons. The area, depth and duration are highly variable and in the altitude range 500–1,000 metres no snow falls in some years. Snowfalls at levels below 500 metres are occasionally experienced in southern Australia, particularly in the foothill areas of Tasmania and Victoria, but falls are usually light and short-lived. In some seasons parts of the eastern uplands above 1,000 metres from Victoria to south-eastern Queensland have been covered with snow for several weeks. In ravines around Mt Kosciuszko (2,228 metres) small areas of snow may persist through summer but there are no permanent snowfields.

### Temperature

*Average temperatures.* Average annual air temperatures as shown in plate 9, page 33 range from 28°C along the Kimberley coast in the extreme north of Western Australia to 4°C in the alpine areas of south-eastern Australia. Although annual temperature may be used for broad comparisons, monthly temperatures are required for detailed analyses.



July is the month with the lowest average temperature in all parts of the continent. The months with the highest average temperature are January or February in the south and December in the north (except in the extreme north and north-west where it is November). The slightly lower temperatures of mid-summer in the north are due to the increase in cloud during the wet season.

*Average monthly maxima.* Maps of average maximum and minimum temperatures for the months of January and July are shown in plates 10–13 inclusive, pages 34–35.


In January, average maximum temperatures exceed 35°C over a vast area of the interior and exceed 40°C over appreciable areas of the north-west. The consistently hottest part of Australia is around Marble Bar, Western Australia (150 kilometres south-east of Port Hedland) where the average is 41°C and daily maxima during summer may exceed 40°C consecutively for several weeks at a time.

The marked gradients of isotherms of maximum temperature in summer in coastal areas, particularly along the south and west coasts, are due to the penetration inland of fresh sea breezes initiated by the sharp temperature discontinuities between the land and sea surfaces. There are also gradients of a complex nature in south-east coastal areas caused primarily by the uplands.



In July a more regular latitudinal distribution of average maxima is evident. Maxima range from 30°C near the north coast to 5°C in the alpine areas of the south-east.



PLATES 8 and 9



PLATES 10 and 11



PLATES 12 and 13

*Extreme maxima.* Temperatures have exceeded 45°C at nearly all inland stations more than 150 kilometres from the coast and at many places on the north-west and south coasts. Temperatures have exceeded 50°C at some inland stations and at a few near the coast. It is noteworthy that Eucla on the south coast has recorded 50.7°C, the highest temperature in Western Australia. This is due to the long trajectory over land of hot north-west winds from the Marble Bar area. Although the highest temperature recorded in Australia was 53.1°C at Cloncurry (Queensland), more stations have exceeded 50°C in western New South Wales than in other areas due to the long land trajectory of hot winds from the north-west interior of the continent.

Extreme maximum temperatures recorded at selected stations, including the highest recorded in each State, are shown in the table below.

**EXTREME MAXIMUM TEMPERATURES**  
(All years to 1976 inclusive)

Station	°C	Station	°C
Western Australia—		New South Wales—	
Eucla . . . . .	50.7	Bourke . . . . .	52.8
Roebourne . . . . .	47.8	White Cliffs . . . . .	51.1
Marble Bar . . . . .	49.2	Walgett . . . . .	50.1
Northern Territory—		Wilcannia . . . . .	50.0
Charlotte Waters (near Finke) . . . . .	48.2	Menindee . . . . .	49.7
South Australia—		Australian Capital Territory—	
Oodnadatta . . . . .	50.7	Canberra . . . . .	42.2
Kyancutta . . . . .	49.3	Victoria—	
Queensland—		Mildura . . . . .	50.8
Cloncurry . . . . .	53.1	Tasmania—	
Winton . . . . .	50.7	Bushy Park . . . . .	40.9
Birdsville . . . . .	50.0	Hobart . . . . .	40.8

*Extreme minima.* The lowest temperatures in Australia have been recorded in the Snowy Mountains, where Charlotte Pass (elevation 1,760 metres) has recorded -22.2°C. Temperatures have fallen below -5°C at most inland places south of the tropics and at some places within a few kilometres of southern coasts. At Eyre, on the south coast of Western Australia, a minimum of -3.9°C has been recorded, and at Swansea, on the east coast of Tasmania, the temperature has fallen as low as -4.4°C.

In the tropics, extreme minima below 0°C have been recorded at many places away from the coasts as far north as Herberton, Queensland (-3.3°C). Even very close to the tropical coastline temperatures have fallen to 0°C, a low recording being -0.8°C for Mackay.

The next table shows extreme minimum temperatures recorded at specified stations, including the lowest recorded in each State.

**EXTREME MINIMUM TEMPERATURES**  
(All years to 1976 inclusive)

Station	°C	Station	°C
Western Australia—		New South Wales—	
Dwellingup . . . . .	-7.0	Charlotte Pass . . . . .	-22.2
Booylgoo . . . . .	-6.7	Kiandra . . . . .	-20.6
Salmon Gums . . . . .	-5.4	Kosciusko Hotel . . . . .	-14.4
Northern Territory—		Cooma . . . . .	-11.2
Alice Springs . . . . .	-7.5	Australian Capital Territory—	
Tempe Downs . . . . .	-6.0	Canberra . . . . .	-10.0
South Australia—		Victoria—	
Yongala . . . . .	-8.2	Mount Hotham . . . . .	-12.8
Kyancutta . . . . .	-7.0	Omeo . . . . .	-11.7
Queensland—		Bairnsdale . . . . .	-7.2
Stanthorpe . . . . .	-11.0	Tasmania—	
Nanango . . . . .	-9.3	Oatlands . . . . .	-12.8
		Bothwell . . . . .	-12.5

*Heat waves.* Periods with a number of successive days having a temperature higher than 40°C are relatively common in summer over parts of Australia. With the exception of the north-west coast of Western Australia, however, most coastal areas rarely experience more than three successive days

of such conditions. The frequency increases inland, and periods of up to ten successive days have been recorded at many inland stations. This figure increases in western Queensland and north-western Western Australia to more than twenty days in places. The central part of the Northern Territory and the Marble Bar-Nullagine area of Western Australia have recorded the most prolonged heat waves.

Heat waves are experienced in the coastal areas from time to time. During 11–14 January 1939, for example, a severe heat wave affected south-eastern Australia: Adelaide had a record of 47.6°C on the 12th, Melbourne a record of 45.6°C on the 13th and Sydney a record of 45.3°C on the 14th.

**Frost.** Frost can cause serious losses in agricultural crops, and numerous climatic studies have been made in Australia relating to specific crops cultivated in local areas. Foley (1945 (i)) made a comprehensive study of the incidence of frost at stations recording minimum temperature. Since Foley's work was published, the number of stations recording minimum temperatures has increased appreciably.

Under calm conditions, overnight temperatures at ground level are often as much as 5°C lower than those measured in the instrument screen (base height 1.1 metre) and differences of 10°C have been recorded. Only a small number of stations measure minima at ground level, the lowest recordings being -15.1°C at Canberra and -14.6°C at Stanthorpe (Queensland). Lower readings may be recorded in alpine areas.

Frost frequency depends on location and orography, and even on minor variations in the contour of the land. The parts of Australia which are most subject to frost are the eastern uplands from north-eastern Victoria to the western Darling Downs in southern Queensland. Most stations in this region experience more than ten nights a month with readings of 0°C (or under) for three to five months of the year. On Tasmania's Central Plateau similar conditions occur for three to six months of the year. Frosts may occur within a few miles of the coasts except on the Northern Territory and most of the north Queensland coasts.




PLATE 14

Regions in which frosts may occur at any time of the year comprise most of Tasmania, large areas of the tablelands of New South Wales, much of inland Victoria, particularly the north-east, and a small part of the extreme south-west of Western Australia. Over most of the interior of the continent, and on the highlands of Queensland as far north as the Atherton Plateau, frosts commence in April and end in September. Minimum temperatures below 0°C are experienced in most of the subtropical interior in June and July.

The length of the frost period for the year is taken as the number of days between the first and last recording of an air temperature of 2°C or less. The median duration of the frost period in days per year is shown in plate 14, page 37.

The median frost period over the continent varies from over 200 days per year in the south-eastern uplands areas south of the Hunter Valley to zero in northern Australia. In the southern regions of the continent the annual frost period generally decreases from about 100 days inland to below 50 days towards the coast. However, there are appreciable spatial variations depending mainly on local orography. In Tasmania the frost period exceeds 300 days on the uplands and decreases to 100 days near the coast.

The table below includes the average annual frequency of minima of 2°C or less for a wide selection of stations, particularly those prone to frosts. These data show the high spatial variability of frost frequency across Australia. The south-eastern alpine areas, as represented by Kiandra (elevation 1,400 metres), have a frequency exceeding 200. At Kalgoorlie the average annual frequency is 27, at Alice Springs 33, Charleville 37, Canberra 105 and Melbourne 19.

#### FROST FREQUENCY

Average annual number of frosty nights (screen minimum  $\leq 2^{\circ}\text{C}$ ) and heavy frosts ( $\leq 0^{\circ}\text{C}$ )

Station	Period of record	Altitude (metres)	Number of frosty nights	Number of heavy frosts
Adelaide (airport)	1955-70	10	8	1
Alice Springs	1940-71	550	33	11
Ballan (near Ballarat)	1944-64	500	63	20
Birdsville	1957-71	40	7	1
Brisbane (Archerfield airport)	1939-49	10	9	3
Canberra	1939-71	570	105	65
Ceduna	1939-71	20	18	5
Charleville	1942-71	290	37	15
Hobart (Risdon)	1957-70	40	25	5
Kalgoorlie	1939-71	360	27	7
Kiandra	1957-69	1,400	226	176
Loch Valley (E of Melbourne)	1943-59	500	101	53
Melbourne (Essendon airport)	1939-71	80	19	4
Mount Gambier	1942-71	60	33	10
Perth (airport)	1944-71	20	5	0
Walgett	1957-71	130	30	7

The next table shows percentiles (20, 50 and 80) of the annual number of frosts at selected stations. The difference between the 20 and 80 percentile figures relative to the 50 percentile (median) shows that there is great variability in the number of frosts at individual stations from year to year.

## FROST VARIABILITY

Annual number of frosty nights ( $\leq 2^{\circ}\text{C}$ ) and heavy frosts ( $\leq 0^{\circ}\text{C}$ ) 20, 50 and 80 percentiles

Station	Period of record	Altitude (metres)	Number of frosty nights			Number of heavy frosts		
			Percentiles			Percentiles		
			20	50	80	20	50	80
Alice Springs	1941-71	550	16	27	37	5	8	14
Bathurst	1957-71	705	83	101	111	51	69	76
Beechworth (SW of Albury)	1957-71	550	51	58	73	16	22	26
Bridgetown	1957-69	155	30	43	53	7	11	19
Canberra	1939-71	570	87	105	116	48	64	75
Charleville	1943-71	290	21	35	45	6	14	19
Dubbo	1957-71	262	39	43	50	10	14	27
Hay	1957-71	93	21	34	37	5	9	13
Kalgoorlie	1942-71	360	15	22	31	2	4	9
Kiandra	1957-68	1,400	206	228	250	163	175	193
Kyancutta	1957-69	58	31	39	40	7	14	20
Mount Gambier	1942-71	60	20	27	34	3	6	13
Mundibindi	1957-69	575	8	11	29	2	3	11
Nhill (near Horsham)	1957-71	129	41	47	58	12	17	26
Oatlands	1957-71	435	85	101	111	38	46	57
Omeo	1957-71	660	115	132	138	59	74	83
Richmond (NW of Sydney)	1953-71	20	23	30	40	6	10	13
Sale	1945-71	5	25	34	45	5	11	17
Swansea	1957-71	8	38	45	61	7	13	19
Wandering (SE of Perth)	1957-69	335	41	57	70	13	25	34
Waratah	1957-71	627	104	117	131	35	44	53
Yongala (E of Port Pirie)	1957-69	515	62	75	90	32	39	52

By convention, a heavy frost is taken as corresponding to a minimum screen temperature of  $0^{\circ}\text{C}$  or less—see the two previous tables. The regions of mainland Australia most prone to heavy frosts are the eastern uplands and adjacent areas extending from Victoria through New South Wales to south-eastern Queensland. Stations above 1,000 metres in altitude in the southern parts of these uplands have more than 100 heavy frosts annually, and in the upland areas below 1,000 metres the annual frequency ranges from 100 to about 20. Over the remainder of southern Queensland, New South Wales and Victoria, although there are great spatial variations, the average annual frequency of heavy frosts typically ranges from about 20 inland to 10 towards the coast. Some places on the coast experience heavy frosts (for example Portland, Victoria, with 3 annually).

In Tasmania, uplands above 1,000 metres have more than 100 heavy frosts annually, and in neighbouring areas the frequency is about 100 decreasing to 20 towards the coasts. Even some coastal stations have a relatively high frequency (Swansea, for example, has 13).

The southern half of Western Australia, the whole of South Australia, and the Alice Springs district of the Northern Territory experience heavy frosts. Differences in annual frequencies between places are great but, in general, the frequency is about 10 inland, decreasing towards the coasts. Some places average more than 20 heavy frosts annually, notably Wandering, Western Australia (21) and Yongala, South Australia (29). At Alice Springs the annual average frequency is 11.

## Humidity

Australia is a dry continent in terms of the water vapour content (humidity) of the air. Humidity is measured at Bureau of Meteorology observational stations by dry and wet bulb thermometers mounted in standard instrument screens. These measurements enable moisture content to be expressed in a number of ways, two of which are vapour pressure and relative humidity.

**Vapour pressure.** Vapour pressure is the pressure exerted by the water vapour in the air and, as such is a measure of the actual amount of water vapour. The amount of water vapour does not normally vary greatly during the day, although afternoon sea breezes at coastal stations may bring in moisture to increase the vapour pressure by amounts up to 5 millibars. The 9 a.m. figure may be taken as an approximation to the mean value for the day. The table on page 41, contains average 9 a.m. vapour pressure figures for selected stations. The average annual figures range from 8.2 millibars at Alice Springs to 25.9 millibars at Darwin and 27.6 millibars at Thursday Island. At the high level station Kiandra (1,400 metres) the average annual figure is 7.3 millibars. Excluding values at Kiandra, monthly averages range from 6.0 millibars at Alice Springs in August to 31.1 millibars at Darwin in January and at both Darwin and Broome in February.

Vapour pressure in association with air temperature has been used as a measure of climatic discomfort as it affects human beings. Comfortable conditions are generally accepted as being within the vapour pressure range 7–17 millibars, with air temperatures in the range  $15\text{--}30^{\circ}\text{C}$ . Above these

limits heat discomfort increases and below the limits cold discomfort increases. The wet bulb temperature may also be used as a simple measure of heat discomfort since discomfort increases as the wet bulb temperature rises above 20°C. Climatic discomfort is treated later in this chapter.

*Relative humidity.* Relative humidity at a given temperature is the ratio (expressed as a percentage) of actual vapour pressure to the saturated vapour pressure at that temperature. The relative humidity at 9 a.m. may be taken as an approximation of the mean relative humidity for the day (24 hours). As a measure of human discomfort this parameter is of limited value because it must be related to the temperature at the time.

The table on page 41 contains average relative humidity (per cent) at 9 a.m. for selected stations. Average annual figures range from 30 per cent at Mundiwindi to 80 per cent at Thursday Island. Monthly averages range from 17 per cent at Mundiwindi in October to 89 per cent at Katanning in June, July and August and at Kiandra in June. In northern Australia the highest relative humidity occurs in the summer rainy season about February and the lowest in the winter dry season about July. Darwin averages 81 per cent in January and February and 62 per cent in July. In most of southern Australia the highest relative humidity is experienced in the winter rainy season about June or July and the lowest in the warmer months. Perth averages 76 per cent in July and 51 per cent in December, January and February. Over the interior, relative humidity is consistently low, although higher averages occur in winter months when temperatures are low. At Alice Springs, October has the lowest average (24 per cent) and June the highest (62 per cent).

The pattern of variation of relative humidity differs from that of vapour pressure, particularly in the south. This is due to the difference in variation of the two parameters with temperature. If the amount of moisture in the air remains constant, vapour pressure decreases slightly with falling temperature, whereas relative humidity increases. Perth, for example, has an average 9 a.m. vapour pressure of 14.8 millibars in January and 10.7 millibars in August; corresponding relative humidity figures are 51 and 71 per cent.

## AVERAGE VAPOUR PRESSURE AT 9 A.M.

(mb)

NOTE. The average monthly and annual figures in this and the next table are derived from the average monthly and annual dry and wet bulb temperatures respectively, using psychrometric formulae. Due to the nature of these formulae annual figures so derived may not equal averages of monthly figures.

Station	Period of record	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year
Adelaide . . .	1868-1971	11.9	12.3	11.7	11.3	10.8	9.9	9.4	9.7	9.9	10.1	10.5	11.1	10.7
Alice Springs . . .	1957-1971	11.9	11.5	10.5	10.1	8.4	7.9	6.5	6.0	6.6	6.8	8.6	9.9	8.2
Armidale . . .	1957-1971	15.1	15.5	14.1	11.7	8.7	7.5	6.3	7.3	8.3	10.1	11.5	13.3	10.2
Brisbane . . .	1887-1950	21.7	22.0	20.9	17.5	14.3	12.1	11.1	11.7	13.8	16.0	18.1	20.1	16.6
Broome . . .	1957-1971	29.4	31.1	29.4	22.4	14.6	14.2	11.8	11.8	15.8	21.7	25.3	28.8	20.8
Canberra . . .	1940-1971	13.1	13.8	12.5	10.3	8.4	7.1	6.6	7.0	8.2	9.7	10.4	11.9	9.9
Carnarvon . . .	1957-1971	21.7	21.9	19.9	16.9	13.8	14.0	11.8	11.6	12.3	13.8	15.9	18.8	15.8
Ceduna . . .	1957-1971	13.8	14.3	14.1	12.0	11.1	9.7	9.4	9.6	10.3	10.3	10.9	12.2	11.1
Charleville . . .	1957-1971	16.7	17.1	15.6	12.5	10.2	9.3	7.8	8.1	8.5	10.5	11.3	14.4	11.3
Cloncurry . . .	1957-1971	19.9	21.2	17.8	13.3	10.4	9.2	7.6	6.9	7.5	9.9	11.8	15.4	12.0
Darwin . . .	1882-1966	31.1	31.1	30.7	27.0	21.8	18.7	17.6	20.6	24.7	27.7	29.3	30.5	25.9
Esperance . . .	1957-1969	16.0	16.7	15.7	14.4	12.7	12.1	11.1	11.1	11.7	12.9	13.9	15.3	13.5
Halls Creek . . .	1957-1971	21.1	21.7	18.5	12.4	10.3	8.2	6.9	6.7	7.5	10.9	13.9	18.0	12.6
Hobart . . .	1894-1970	11.0	11.7	11.0	10.0	8.8	7.9	7.6	7.9	8.3	9.1	9.6	10.6	9.5
Kalgoorlie . . .	1957-1971	12.9	14.0	13.1	11.8	10.3	10.1	8.9	8.8	9.1	9.6	10.5	11.7	10.7
Katanning . . .	1957-1972	13.2	13.9	13.2	12.5	11.0	10.5	9.3	9.7	10.2	9.4	10.6	11.5	11.1
Kiandra . . .	1957-1972	11.1	11.3	10.3	7.6	5.9	5.4	4.7	5.2	5.5	7.3	8.1	10.3	7.3
Marble Bar . . .	1957-1971	20.4	20.8	17.8	12.6	9.5	10.3	7.8	7.6	7.8	9.1	11.0	15.0	11.9
Melbourne . . .	1907-1971	13.1	14.1	13.3	11.7	10.3	9.3	8.9	9.1	9.5	10.5	11.3	12.5	11.1
Mildura . . .	1957-1971	13.6	13.7	13.1	11.7	10.3	9.0	8.7	9.0	9.9	10.4	10.8	11.9	10.8
Mundawindi . . .	1957-1972	13.1	14.4	11.8	10.6	8.5	8.8	7.2	6.8	6.7	6.4	8.2	10.2	8.9
Perth . . .	1911-1940	14.8	14.7	14.7	13.4	12.4	11.4	10.9	10.7	11.6	11.7	12.7	13.9	12.7
Sydney . . .	1876-1971	18.8	19.2	18.3	15.0	11.9	10.2	9.6	9.5	11.3	13.0	15.0	17.6	13.6
Thursday Island . . .	1957-1971	30.2	30.4	30.3	29.0	28.0	25.8	24.1	24.5	24.7	26.1	28.0	29.6	27.6
Townsville . . .	1957-1971	26.1	27.3	25.4	22.1	18.2	15.3	14.1	15.7	16.7	19.7	22.9	24.6	20.3

## AVERAGE RELATIVE HUMIDITY AT 9 A.M.

(per cent)

Station	Period of record	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Year
Adelaide . . .	1868-1971	41	43	47	56	67	75	76	70	60	51	45	40	56
Alice Springs . . .	1957-1971	30	32	35	44	54	62	56	43	33	24	25	27	35
Armidale . . .	1957-1971	63	68	70	73	77	80	74	73	63	56	54	57	66
Brisbane . . .	1887-1971	66	69	71	71	71	72	70	67	63	60	59	61	67
Broome . . .	1957-1971	69	74	69	56	46	52	48	43	46	54	58	64	58
Canberra . . .	1940-1971	58	65	67	73	83	85	83	78	72	66	57	56	69
Carnarvon . . .	1957-1971	62	59	59	59	60	73	68	63	55	53	55	59	61
Ceduna . . .	1957-1971	49	54	60	61	75	77	80	74	63	49	45	48	59
Charleville . . .	1957-1971	47	49	52	52	62	71	65	55	42	38	34	41	48
Cloncurry . . .	1957-1971	48	54	49	42	45	49	43	34	28	27	28	35	40
Darwin . . .	1882-1971	81	81	80	72	65	63	62	66	68	68	70	75	71
Esperance . . .	1957-1969	61	67	66	71	75	82	83	77	71	63	62	62	69
Halls Creek . . .	1957-1971	48	51	44	31	34	33	29	24	20	23	28	38	34
Hobart . . .	1894-1971	58	62	65	70	75	78	78	73	66	62	58	58	67
Kalgoorlie . . .	1957-1971	43	50	52	58	65	75	74	65	55	46	42	42	54
Katanning . . .	1957-1972	58	65	68	78	78	89	89	89	83	54	51	48	69
Kiandra . . .	1957-1972	63	68	72	75	85	89	88	87	71	64	58	64	71
Marble Bar . . .	1957-1971	40	44	38	31	32	43	36	30	24	21	22	28	32
Melbourne . . .	1907-1971	60	63	66	72	79	83	81	75	68	63	60	60	69
Mildura . . .	1957-1971	49	52	59	69	82	87	88	81	69	56	48	48	63
Mundawindi . . .	1957-1972	28	35	30	35	40	50	46	37	28	17	19	21	30
Perth . . .	1911-1940	51	51	57	61	70	75	76	71	66	60	52	51	62
Sydney . . .	1876-1971	68	70	74	74	75	76	74	68	66	62	62	64	69
Thursday Island . . .	1957-1971	84	86	85	81	82	80	79	79	75	73	73	77	80
Townsville . . .	1957-1971	69	75	73	68	66	66	64	63	56	58	62	64	65

**Sunshine, cloud and fog**

*Sunshine.* Sunshine as treated here refers to bright or direct sunshine. Australia receives relatively large amounts of sunshine although seasonal cloud formations have a notable effect on its spatial and temporal distribution. Cloud cover reduces both incoming and outgoing radiation and thus affects sunshine, air temperature and other climatic elements at the earth's surface. Sunshine amounts at Australian capitals are included in the climatic tables, pages 54-61.



Average daily sunshine (hours) in January and July based on all available data to August 1974 is shown in plates 15 and 16, page 43. In areas where there is a sparsity of data, estimates of sunshine derived from cloud data were used. Most of the continent receives more than 3,000 hours of sunshine a year, or nearly 70 per cent of the total possible. In central Australia and the mid-west coast of Western Australia totals slightly in excess of 3,500 hours occur. Totals of less than 1,750 hours occur on the west coast and highlands of Tasmania; this amount is only 40 per cent of the total possible per year (about 4,380 hours).

In southern Australia generally the duration of sunshine is greatest about December when the sun is at its highest elevation and lowest in June when the sun is lowest. In northern Australia sunshine is generally greatest about August-October prior to the wet season and least about January-March during the wet season. The table below gives the 20, 50 and 80 percentiles of daily bright sunshine for the months of June and December at selected stations. These values give an indication of the variability of daily sunshine hours. Perth, for example, has a high variability of daily sunshine hours in the wet month of June (160 per cent) and a low variability in the dry month of December (30 per cent). Darwin has a low variability in the dry season month of June (15 per cent) and a high variability in the wet season month of December (85 per cent).

**BRIGHT SUNSHINE, VARIABILITY OF DAILY HOURS, JUNE AND DECEMBER**  
(20, 50 and 80 percentile values)

Station	Period of record	June			December		
		Percentile			Percentile		
		20	50	80	20	50	80
Adelaide . . .	1955-71	1.0	3.5	7.5	4.0	9.0	12.5
Alice Springs . . .	1954-71	5.5	9.5	10.0	6.5	11.0	12.5
Brisbane . . .	1951-71	2.5	8.0	9.5	4.0	8.5	11.5
Canberra . . .	1957-71	2.0	5.0	7.0	4.0	9.5	12.0
Darwin . . .	1951-71	9.0	10.0	10.5	3.5	7.5	10.0
Hobart . . .	1955-71	0.5	3.0	6.0	2.5	7.0	10.5
Melbourne . . .	1955-70	0.5	2.5	6.0	3.0	7.5	11.5
Perth . . .	1945-71	1.0	4.0	7.5	8.5	11.0	12.0
Sydney . . .	1955-71	0.5	6.0	8.0	1.5	7.5	11.0
Townsville . . .	1957-71	4.5	9.0	10.0	5.0	9.5	11.0

*Cloud.* Seasonal changes in cloudiness vary with the distribution of rainfall. In the southern parts of the continent, particularly in the coastal and low lying areas, the winter months are generally more cloudy than the summer months. This is due to the formation of extensive areas of stratiform cloud and fog during the colder months, when the structure of the lower layers of the atmosphere favours the physical processes resulting in this type of cloud. Particularly strong seasonal variability of cloud cover exists in northern Australia where skies are clouded during the summer wet season and mainly cloudless during the winter dry season. Cloud coverage is greater near coasts and on the windward slopes of the eastern uplands of Australia and less over the dry interior.



The average monthly cloud amounts at Australian capitals are included in the climatic tables on pages 54–61. Darwin has the least average daily coverage of 3.2 eighths and Hobart the highest daily average of 5.0 eighths. The highest daily average for any month occurs at Darwin (5.9 eighths for January) and the lowest average daily is also at Darwin (1.1 eighths for August).

**Fog.** The formation of fog depends on the occurrence of favourable meteorological elements—mainly temperature, humidity, wind and cloud cover. The nature of the local terrain is important for the development of fog and there is a tendency for this phenomenon to persist in valleys and hollows. The incidence of fog may vary significantly over distances as short as one kilometre.

Fog in Australia tends to be greater in the south than the north, although parts of the east coastal areas are relatively fog prone even in the tropics. Incidence is much greater in the colder months, particularly in the eastern uplands. Fog may persist during the day but rarely until the afternoon over the interior. The highest fog incidence at a capital city is at Canberra which has an average of 46 days per year on which fog occurs, 28 of which are in the period May to August. Brisbane averages 22 days of fog per year, 17 of which occur between April and September. Darwin averages only 3 days per year, June to September.

### Global radiation

Global (short wave) radiation includes that radiation energy reaching the ground directly from the sun and that received indirectly from the sky, scattered downwards by clouds, dust particles, etc.

Plates 17 and 18, page 45, show the average global radiation for the months of January and July. The table below shows the variability of daily global radiation for June and December (1968–72) at selected stations.

#### GLOBAL RADIATION: VARIABILITY OF DAILY AMOUNTS FOR JUNE AND DECEMBER

(mWh . cm<sup>-2</sup>)

(20, 50 and 80 percentile values in milliwatt hours per square centimetre (1964–68))



Station	June			December		
	Percentiles			Percentiles		
	20	50	80	20	50	80
Alice Springs . . .	360	450	480	580	760	810
Darwin . . . . .	520	570	590	440	570	620
Melbourne . . . .	130	190	240	470	640	780
Perth . . . . .	180	260	330	770	870	910
Townsville . . . .	360	490	510	550	710	760
Williamtown . . . .	210	270	330	490	650	780

A high correlation exists between daily global radiation (plates 17 and 18, page 45) and daily hours of sunshine (plates 15 and 16, page 43). On the north-west coast around Port Hedland, where average daily global radiation is the highest for Australia (640 milliwatt hours), average daily sunshine is also highest, being approximately 10 hours. Sunshine is more dependent on variations in cloud coverage than is global radiation, since the latter includes diffuse radiation from the sky as well as direct radiation from the sun. An example is Darwin where in the dry month of July sunshine approaches twice that of the wet (cloudy) month of January but global radiation figures for the two months are comparable.

### Evaporation

Evaporation is determined by measuring the amount of water evaporated from a free water surface exposed in a pan. Evaporation from a free water surface depends on a number of climatic elements, mainly temperature, humidity and wind. Evaporation data are useful in water conservation studies and in estimating potential evapotranspiration for irrigation and plant growth studies. In Australia, where surface water storage is vital over large areas, evaporation is a highly significant element.

Average annual Class A pan evaporation is mapped in plate 19, page 46, which shows a variation from 900 millimetres in southwest Tasmania to 4,500 millimetres in the dry interior of Western Australia with about 75 per cent of the continent exceeding 2,500 millimetres. In about 75 per cent of the continent, comprising most inland areas, rainfall does not exceed evaporation loss from a free water surface in any month of the year. In the central and north-west parts of the continent the annual evaporation exceeds ten times the rainfall.






PLATE 19

### Winds

The mid-latitude anticyclones are the chief determinants of Australia's two main prevailing wind streams. In relation to the west-east axes of the anticyclones these streams are easterly to the north and westerly to the south. The cycles of development, motion and decay of low pressure systems to the north and south of the anticyclones result in diversity of wind flow patterns. Wind variations are greatest around the coasts where diurnal land and sea breeze effects are important.

Wind roses for the months of January and July at 9 a.m. and 3 p.m. at selected stations are shown in Plates 20–23 inclusive, pages 47–48. The wind roses show the percentage frequency of direction (eight points of compass) and speed ranges of winds.



Orography affects the prevailing wind pattern in various ways such as the channelling of winds through valleys, deflection by mountains and cold air drainage from highland areas. An example of this channelling is the high frequency of north-west winds at Hobart caused by the north-west south-east orientation of the Derwent River Valley.

Average wind speeds and prevailing directions at Australian capitals are included in the climatic tables on pages 54–61. Perth is the windiest capital with an average wind speed of 15.6 kilometres per hour; Canberra is the least windy with an average speed of 5.8 kilometres per hour.



The highest wind speeds and wind gusts recorded in Australia have been associated with tropical cyclones. The highest recorded gust was 246 kilometres per hour during a cyclone at Onslow, Western Australia in 1975 and gusts reaching 200 kilometres per hour have been recorded on several occasions in northern Australia with cyclone visitations. The highest gusts recorded at Australian capitals were 217 kilometres per hour at Darwin and 156 kilometres per hour at Perth.

Estimates of the extreme wind gust expected in a given return period\* have been derived for places throughout Australia (Whittingham, 1964). On this basis, for example, Darwin would have an extreme gust for a return period of 10 years of 140 kilometres per hour, Melbourne 135 and Perth 130.

\*Return period is the average period between successive occurrences equal to, or greater than, a given speed. For example the extreme wind gust for a return period of 10 years can be expected to occur once in 10 years on the average.



PLATES 20 and 21



PLATES 22 and 23

### Floods

Widespread flood rainfall may occur anywhere in Australia but it has a higher incidence in the north and in the eastern coastal areas. It is most economically damaging along the shorter streams flowing from the eastern uplands eastward to the seaboard of Queensland and New South Wales. These flood rains are notably destructive in the more densely populated coastal river valleys of New South Wales—the Tweed, Richmond, Clarence, Macleay, Hunter and Nepean-Hawkesbury—all of which experience relatively frequent flooding. Although chiefly summer rains, they may occur in any season.

The great Fitzroy and Burdekin river basins of Queensland receive flood rains during the summer wet season. Much of the run-off due to heavy rain in north Queensland west of the eastern uplands flows southward through the normally dry channels of the network of rivers draining the interior lowlands into Lake Eyre. This widespread rain may cause floods over an extensive area, but it soon seeps away or evaporates, occasionally reaching the lake in quantity. The Condamine and other northern tributaries of the Darling also carry large volumes of water from flood rains south through western New South Wales to the Murray and flooding occurs along their courses at times.

Flood rains occur at irregular intervals in the Murray-Murrumbidgee system of New South Wales and Victoria, the coastal streams of southern Victoria and the north coast streams of Tasmania.

### Droughts

Drought, in general terms, refers to an acute water shortage. This is normally due to rainfall deficiency but with other parameters contributing to the actual water availability. The best single measure of water availability in Australia is rainfall, although parameters such as evaporation and soil moisture are significant, or even dominant, in some situations.

Droughts have severe economic effects in Australia and during the years 1864–1973 inclusive there have been at least eight major droughts affecting the greater part of Australia and at least seven other droughts of lesser severity affecting extensive areas (Foley 1957 (ii)). The droughts of 1895–1903 and 1958–68 were probably the most disastrous in their effects on primary industry.

Gibbs and Maher (1967), having defined a drought year at a certain station as one with the year's rainfall in the first decile range, concluded that the occurrence of areas in the first decile range on annual decile maps for the period 1885–1965 corresponded rather well with drought areas discussed by Foley (1957).

One method of assessing the incidence of rainfall deficiency is the analysis of the distribution of annual rainfalls less than the median (Gaffney 1975). The range between the 50 percentile (median) and the 10 percentile gives a measure of the variation in magnitude of annual rainfalls less than the median. The ratio of this range to the 30 percentile value may be used as an index of rainfall deficiency incidence or drought incidence, i.e.:

$$\text{Index of drought incidence} = \left\{ \frac{50 - 10}{30} \right\} \text{ percentile}$$

For example, the indexes for Onslow (north-west coast of Western Australia) and similarly, for Cape Otway (south coast of Victoria) are derived thus:

$$\begin{aligned} \text{Index for Onslow} &= \left\{ \frac{222 - 64}{145} \right\} \text{ mm} = 1.09 \\ \text{Index for Cape Otway} &= \left\{ \frac{865 - 716}{801} \right\} \text{ mm} = 0.19 \end{aligned}$$

Plate 24, page 50, shows the distribution of the index of drought incidence over Australia. The intrusions of high index values from the interior to the central coast of Queensland and across western New South Wales are noteworthy. The extreme values on the north-west coast of Western Australia are among the highest in Australia (e.g. Onslow 1.09) due to the dependence of the rainfall on random cyclone tracks.

The Bureau of Meteorology commenced the issue of *Drought Reviews* in June 1965. These reviews provide a summary of serious rainfall deficiencies and are issued monthly when serious or severe deficiencies exist in any of the rainfall districts. The deficiency criteria are based on monthly rainfall decile analyses. A review of droughts in Australia to 1968 is included in Year Book No. 54, 1968. Summaries of subsequent drought periods may be obtained from the *Drought Reviews*.




PLATE 24

### Climatic discomfort

In Australia climatic discomfort is significant in most areas. During the summer half of the year (November–April) prolonged high temperatures and humidity around the northern coasts and high temperatures over the inland cause physical stress. In winter, low temperatures and strong cold winds over the interior and southern areas can be severe for relatively short periods. However, cold stress does not cause prolonged physical hardship in Australia at altitudes lower than 1,000 metres, that is, over more than 99 per cent of the continent.

The climatic variables determining physical discomfort are primarily air temperature, vapour pressure and wind. The complete assessment of physical discomfort also requires analyses of such parameters as thermal conductivity of clothing, vapour pressure at the skin and the metabolic heat rate arising from activity of the human body. The cooling system of the human body depends on evaporation of moisture to keep body temperature from rising to lethal levels as air temperature rises. Defining criteria of discomfort is difficult because personal reactions to the weather differ greatly according to a number of variables including health, age, clothing, occupation and acclimatisation (Ashton 1964). However, climatic strain has been measured experimentally and discomfort indexes based on the average response of subjects under specified conditions have been derived.

**Effective Temperature.** The effective temperature with respect to any environmental combination of temperature, humidity and wind is defined as the temperature of still, saturated air in which a normally clothed sedentary worker would feel the same level of comfort or discomfort.

Environment studies carried out at the research laboratories of the American Society of Heating, Refrigerating and Air Conditioning Engineers established values of effective temperature corresponding to various combinations of temperature, humidity and air movement. The results were published as a series of research reports commencing in 1923, and have been widely used to measure climatic discomfort (see 1960 report of the Society).

Normally clothed sedentary workers are mostly comfortable within a range of effective temperatures between 15°C and 27°C (air movement 5–8 metres per minute). At effective temperatures greater than 27°C, the majority of people feel heat discomfort and when less than 15°C they feel cold discomfort.

The table below contains the annual average frequency of effective temperature at 3 p.m. within specified limits at selected stations. The figures provide comparisons of daily occurrence of afternoon discomfort for the given environmental conditions.

#### CLIMATIC DISCOMFORT: EFFECTIVE TEMPERATURE

Annual average frequency of days when effective temperature at 3 p.m. is lower than 15°C (cold discomfort), within 15–27°C (comfort), and higher than 27°C (heat discomfort). Indoors, normally clothed sedentary workers, air movement 5–8 metres per minute.

Station	Period of record	Average days per year		
		Less than 15°C	15–27°C	Greater than 27°C
Adelaide . . . . .	1955–72	128	234	3
Albury . . . . .	1962–71	141	220	4
Alice Springs . . . . .	1955–67	39	300	26
Brisbane . . . . .	1951–70	6	356	3
Broome . . . . .	1941–71	0	225	140
Canberra . . . . .	1940–72	172	192	1
Carnarvon . . . . .	1945–72	1	345	19
Ceduna . . . . .	1955–71	77	279	9
Charleville . . . . .	1942–72	28	316	21
Cloncurry . . . . .	1940–72	1	268	96
Darwin . . . . .	1955–69	0	225	140
Hobart . . . . .	1944–67	239	126	0
Kalgoorlie . . . . .	1940–72	66	281	18
Marble Bar . . . . .	1957–71	0	220	145
Melbourne . . . . .	1955–71	155	207	3
Mildura . . . . .	1946–72	95	258	12
Perth . . . . .	1944–71	57	302	6
Rockhampton . . . . .	1940–72	2	337	26
Sydney . . . . .	1955–72	69	295	1
Townsville . . . . .	1941–69	0	333	32
Woomera . . . . .	1954–72	73	279	13

Heat discomfort is greatest in the north-west, where Marble Bar averages 145 days of high heat discomfort annually, and least in the south-east, where Hobart has only one day every five years. Cold discomfort is least in the north, where Townsville has one day of cold discomfort in ten years, and greatest in the south-east, where Hobart has 239 days annually when the effective temperature is sufficiently low to cause discomfort. By the suitable choice of clothing discomfort can be decreased significantly on cold days. On cold days also, workers tend to take opportunities to move around, thus increasing metabolic heat rates.

Effective temperature is a useful index but its application is limited because available criteria relate only to indoor workers in sedentary occupations. Furthermore, at lower air temperatures the effective temperature gives excessive weight to humidity.

*Relative strain index.* The relative strain index derived by Lee and Henschel (1963) has been applied in Australia to measure heat discomfort (Hounam, 1969, Gaffney 1973). The results obtained with Australian data are useful for purposes of comparison but interpretation of the actual results is tentative until empirical environmental studies are carried out in this region. In addition to temperature, humidity and air movement the relative strain index has facilities for incorporation of metabolic heat rate, net radiation and insulation of clothing. It has the advantage of being applicable to manual workers under shelter and expending energy at various metabolic heat rates.

The discomfort map plate 25, page 53, shows the average number of days per year when the relative strain index exceeds 0.3 discomfort level at 3 p.m. assuming standard conditions as defined. Maximum discomfort generally occurs around 3 p.m. on days of high temperature.

A notable feature is the lower frequency of days of discomfort in Queensland coastal areas in comparison with the northern coastal areas of Western Australia. This is due to the onshore winds prevailing on the Queensland coast and the cooling effect of the adjacent eastern uplands. Lower frequencies on the Atherton Plateau in the tropics near Cairns show the advantage of altitude. Relatively low heat discomfort frequencies are evident in upland and coastal areas of south-east Australia. Tasmania is entirely in the zone of least discomfort, experiencing on the average less than one day of heat discomfort per year. In Western Australia most of the Kimberley region in the north lies in the highest discomfort zone with the frequencies decreasing southwards to a strip of lowest discomfort towards the south-west coast. A steep gradient of discomfort frequency on the west coast shows the moderating effect of sea breezes.

The average annual frequency of days when the relative strain index at 3 p.m. exceeds specified discomfort levels is shown in the table below. The Sydney frequencies were derived from observations at the Regional Office of the Bureau of Meteorology, which is representative of eastern coastal suburbs; frequencies are higher in western suburbs. The Melbourne frequencies were derived from observations at the Bureau's Regional Office, which may be taken as fairly representative of inner northern and eastern suburbs; frequencies are lower in bayside suburbs. Similarly, in other capital city areas significant variations occur with distance from the coast.

#### HEAT DISCOMFORT

Average number of days per year when relative strain index (RSI) at 3 p.m. exceeds 0.3 (discomfort) and 0.4 (high discomfort) under standard conditions (indoors, manual activities, light clothing, air movement 60 metres per minute).

Station	Period of record	Greater than	
		0.3 RSI	0.4 RSI
Adelaide . . . . .	1955-72	7	1
Albury . . . . .	1962-71	8	1
Alice Springs . . . . .	1955-67	50	4
Brisbane . . . . .	1951-69	6	<1
Broome . . . . .	1940-72	155	48
Canberra . . . . .	1940-72	2	<1
Carnarvon . . . . .	1945-72	23	3
Ceduna . . . . .	1955-71	16	3
Charleville . . . . .	1942-72	42	3
Cloncurry . . . . .	1940-72	126	28
Darwin . . . . .	1955-69	165	23
Hobart . . . . .	1944-67	<1	<1
Kalgoorlie . . . . .	1939-72	30	5
Marble Bar . . . . .	1957-71	173	69
Melbourne . . . . .	1955-71	6	1
Mildura . . . . .	1946-72	19	3
Perth . . . . .	1944-72	12	1
Rockhampton . . . . .	1940-72	33	5
Sydney . . . . .	1955-72	2	<1
Townsville . . . . .	1941-69	36	4
Woomera . . . . .	1954-72	25	3

At inland places, relatively low night temperatures have recuperative effects after hot days. Marble Bar, Western Australia (150 km south-east of Port Hedland) for example, has median night minimum temperatures 5-10° C lower than Darwin, except in December-February. Even in this latter period, although median minima at both stations are around 25° C, Marble Bar has median vapour pressures and relative humidities much lower than Darwin (by 10 millibars and 30 per cent respectively).

Acclimatised people would suffer discomfort less frequently than shown by the relative strain index figures. For example, Australians living in the north evidently experience less discomfort at high air temperatures than those in the south, if humidities are comparable.

Both direction and speed of prevailing winds are significant for the ventilation of buildings. In the tropics, for instance, windward slopes allow optimal air movement enabling more comfortable ventilation to be obtained. Regular sea breezes such as those experienced at Perth reduce discomfort although on some days their full benefit may not be experienced until after 3 p.m.

#### Climatic data for capital cities

The averages for a number of elements determined from long-period observations at the Australian capitals to 1976 inclusive, are given in the following pages. Extremes generally cover all available data to 1976 inclusive, whereas averages may only refer to present sites.




PLATE 25

## CLIMATIC DATA: PERTH, WESTERN AUSTRALIA

(Lat. 31° 57' S., Long. 115° 52' E. Height above M.S.L. 15 metres)

## BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 22 metres)					Mean amt evapo- ration (mm)	No. days thun- der	Mean daily amt clouds 9 a.m., 3 p.m., 9 p.m. (a)	No. clear days
		Aver- age (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction					
					9 a.m.	3 p.m.				
No. of years of record .	91	30(b)	76	61	30(b)	30(b)	9	79	30(b)	30(b)
January . . . . .	1,012.6	17.5	48.2 26/76*	81	E	SSW	280	0.9	2.3	14
February . . . . .	1,013.0	17.2	40.8 4/73	87	ENE	SSW	241	0.7	2.5	13
March . . . . .	1,015.2	16.2	51.9 28/75	113	E	SSW	214	0.7	2.8	12
April . . . . .	1,017.9	13.7	50.7 25/100	101	ENE	SSW	124	0.9	3.4	9
May . . . . .	1,017.9	13.5	44.5 8/73	119	NE	WSW	83	1.8	4.3	6
June . . . . .	1,017.5	13.5	48.6 17/27	129	N	NW	59	1.8	4.7	6
July . . . . .	1,018.8	14.2	53.9 20/26	137	NNE	W	58	1.5	4.5	6
August . . . . .	1,018.8	15.1	51.3 15/03	156	N	WNW	75	1.3	4.5	6
September . . . . .	1,018.4	15.1	45.9 11/05	109	ENE	SSW	105	0.7	3.9	8
October . . . . .	1,017.0	16.1	43.0 6/16	105	SE	SW	158	0.8	3.8	8
November . . . . .	1,015.5	17.2	48.2 26/75*	101	E	SW	205	0.8	3.1	9
December . . . . .	1,013.4	17.7	44.5 24/75	103	E	SSW	241	0.9	2.6	13
Year { Totals . . . . .	1,016.3	15.6	..	..	E	SSW	1,843	12.8	3.5	108
Year { Averages . . . . .	..	..	..	..	..	..	..	..	..	..
Year { Extremes . . . . .	..	..	53.9	156	..	..	..	..	..	..
			20/7/26							

(a) Scale 0-8. (b) Standard thirty years normal (1911-1940).

## TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (°Celsius)			Extreme air temperature (°Celsius)		Extreme temperature (°Celsius)		Mean daily hours sun-shine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record .	79	79	79	80	80	63(a)	78	78
January . . . . .	29.5	17.6	23.5	43.7 29/56	9.2 20/25	80.7 22/14	4.2 20/25	10.5
February . . . . .	29.8	17.8	23.7	44.6 8/33	8.7 1/02	78.7 4/34	4.3 1/13	10.0
March . . . . .	27.8	16.5	22.2	41.3 14/22	7.7 8/03	75.0 19/18	2.6 (b)	8.9
April . . . . .	24.4	14.0	19.2	37.6 9/10	4.1 20/14	69.4 8/16	-0.7 26/60	7.2
May . . . . .	20.6	11.5	16.1	32.4 2/07	1.3 11/14	63.3 4/25	-3.9 31/64	5.9
June . . . . .	18.1	9.9	14.1	28.1 5/75	1.6 22/55	57.5 9/14	-3.4 27/46	4.8
July . . . . .	17.3	9.0	13.2	26.3 17/76	1.2 7/16	56.2 13/15	-3.8 30/20	5.3
August . . . . .	17.9	9.1	13.5	27.8 21/40	1.9 31/08	62.3 29/21	-3.0 18/66	6.2
September . . . . .	19.4	10.1	14.8	32.7 30/18	2.6 6/56	67.5 29/16	-2.7 (c)	7.2
October . . . . .	21.2	11.4	16.3	37.3 29/67	4.2 6/68	71.8 19/54	-1.2 16/31	8.3
November . . . . .	24.5	13.8	19.2	40.3 24/13	5.6 1/04	75.0 30/25	-1.1 6/71	9.7
December . . . . .	27.3	16.1	21.7	42.3 31/68	8.6 29/57	76.0 11/27	3.3 29/57	10.8
Year { Averages . . . . .	23.2	13.1	18.2	..	..	..	..	7.9
Year { Extremes . . . . .	..	..	..	44.6	1.2	80.7	-3.9	..
				8/2/33	7/7/16	22/1/14	31/5/64	

(a) Records discontinued 1963. (b) 8/1903 and 16/1967. (c) 8/1952 and 6/1956.

## HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)					Fog Mean No. days
		Mean	Highest mean	Lowest mean	Mean No. of days of rain monthly	Greatest monthly	Least monthly	Greatest in one day		
No. of years of record .	30(a)	30(a)	79	79	100	100	100	100	100	79
January . . . . .	14.8	51	63	41	8	3	55 1879	Nil (b)	44 27/79*	0.2
February . . . . .	14.7	51	65	43	11	3	166 1955	Nil (b)	87 17/55	0.3
March . . . . .	14.7	57	66	46	20	4	145 1934	Nil (b)	77 9/34	0.6
April . . . . .	13.4	61	75	51	46	8	149 1926	Nil 1920	67 30/04	0.9
May . . . . .	12.4	70	81	60	125	14	308 1879	14 1964	76 17/42	1.3
June . . . . .	11.4	75	85	68	185	17	476 1945	55 1877	99 10/20	1.4
July . . . . .	10.9	76	88	69	175	18	425 1958	61 1876	76 4/91*	1.6
August . . . . .	10.7	71	83	62	138	18	318 1945	12 1902	74 14/45	1.0
September . . . . .	11.6	66	75	58	81	14	199 1923	9 1916	47 18/66	0.3
October . . . . .	11.7	60	75	52	55	11	200 1890	1 1969	50 4/67	0.4
November . . . . .	12.7	52	66	41	21	6	71 1916	Nil 1891	39 29/56	0.2
December . . . . .	13.9	51	63	39	14	4	81 1951	Nil (b)	47 3/51	0.2
Year { Totals . . . . .	..	..	..	..	879	120	..	..	..	8.1
Year { Averages . . . . .	12.7	62	..	..	..	..	..	..	..	..
Year { Extremes . . . . .	..	..	88	39	..	..	476	Nil (b)	99	..
							6/1945		10/620	

(a) Standard thirty years normal (1911-1940). (b) Various years.

Figures such as 26/76, 29/56, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk(\*) relate to nineteenth century.

## CLIMATIC DATA: DARWIN, NORTHERN TERRITORY

(Lat. 12° 28' S., Long. 130° 51' E. Height above M.S.L. 30 metres)

## BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 36 metres)				Mean amt evaporation (mm)	No. days thunder	Mean daily amt clouds	
		Average (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction			9 a.m., 3 p.m., 9 p.m.	No. clear days
No. of years of record .	90	20	..	22(b)	..	8	35	35	35
January . . . . .	1,006.2	9.3	..	106	W	NW	225	12.9	5.9
February . . . . .	1,006.3	10.6	..	101	W	NW	187	10.2	5.8
March . . . . .	1,007.2	7.5	..	157	W	NW	190	10.6	5.2
April . . . . .	1,009.3	8.8	..	67	SE	NW	218	4.0	2.9
May . . . . .	1,010.9	9.6	..	62	SE	E	223	0.5	2.0
June . . . . .	1,012.2	10.1	..	64	SE	E	206	0.0	1.4
July . . . . .	1,012.8	8.9	..	62	SE	E	229	0.0	1.3
August . . . . .	1,012.6	8.6	..	72	SE	NW	238	0.0	1.1
September . . . . .	1,011.7	8.6	..	64	ENE	NW	270	1.0	1.8
October . . . . .	1,010.5	9.8	..	85	NE	NW	285	5.3	2.7
November . . . . .	1,008.7	8.6	..	117	NW	NW	260	11.8	3.9
December . . . . .	1,006.9	9.8	..	217	NW	NW	240	14.2	4.9
Year { Totals . . . . .	..	..	..	..	..	2,773	70.5	..	121
Year { Averages . . . . .	1,009.6	9.2	..	..	SE	NW	..	..	3.2
Year { Extremes . . . . .	..	..	..	217	..	..	..	..	..

(a) Scale 0-8. (b) Several incomplete years.

## TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (°Celsius)			Extreme air temperature (°Celsius)		Extreme temperature (°Celsius)		Mean daily hours sun-shine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record .	90	90	90	92(a)	92(a)	26(b)	..	21
January . . . . .	32.2	25.0	28.6	37.8 2/82*	20.0 20/92*	75.6 26/42	..	5.9
February . . . . .	31.9	24.8	28.4	38.3 20/87*	17.2 25/49	73.2 (c)	..	5.9
March . . . . .	32.4	24.8	28.6	38.9 (d)	19.2 31/45	74.3 23/38	..	6.8
April . . . . .	33.1	24.2	28.7	40.0 7/83*	16.0 11/43	72.8 1/38	..	8.6
May . . . . .	32.3	22.4	27.4	39.1 8/84*	(e) 14.2 28/67	71.2 5/20	..	9.3
June . . . . .	30.9	20.4	25.7	39.0 17/37*	12.1 23/63	68.5 2/16	..	9.7
July . . . . .	30.4	19.6	25.1	36.7 17/88*	10.4 29/42	68.9 28/17	..	9.8
August . . . . .	31.4	20.8	26.1	37.0 30/71*	13.6 11/63	69.1 28/16	..	10.4
September . . . . .	32.7	23.2	27.9	38.9 20/82*	16.7 9/63	69.5 (f)	..	10.0
October . . . . .	33.6	25.0	29.3	40.5 17/92*	19.4 8/66	71.4 30/38	..	9.5
November . . . . .	33.8	25.3	29.6	39.6 9/84*	19.3 4/50	77.0 14/37	..	8.6
December . . . . .	33.2	25.3	29.3	38.9 20/82*	18.3 4/60	76.2 26/23	..	7.1
Year { Averages . . . . .	32.3	23.3	27.9	..	..	..	..	8.5
Year { Extremes . . . . .	..	..	..	40.5	10.4	77.0	..	..
				17/10/1892	29/7/1942	14/11/37		

(a) Years 1882-1941 at Post Office, 1942-1966 at Aerodrome; 1967-1971 at Regional office; sites not strictly comparable. (b) Records discontinued 1942. (c) 5/1938 and 23/1938. (d) 26/1883 and 27/1883. (e) Recorded at Darwin Aerodrome. All other Statistics from 1967 to 1971 at Regional Office. (f) 28/1916 and 3/1921.

## HUMIDITY, RAINFALL, AND FOG

Month	Vapour pres- sure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)					Fog Mean No. days
		Mean	Highest mean	Lowest mean	Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day	
No. of years of record .	85(a)	90	57(b)	57(b)	86(c)	74	107(d)	107(d)	107(d)	35
January . . . . .	31.1	81	89	69	391	19	746 1974	68 1906	296 7/97*	0.0
February . . . . .	31.1	81	88	71	330	18	815 1969	13 1931	279 18/55	0.0
March . . . . .	30.7	80	84	69	260	17	1013 1977	21 1911	241 16/77	0.0
April . . . . .	27.0	72	80	60	103	8	603 1891	Nil 1950	158 4/59	0.0
May . . . . .	21.8	65	76	49	14	1	299 1968	Nil (e)	56 6/22	0.0
June . . . . .	18.7	63	75	52	3	0	76 1973	Nil (e)	36 0/02	0.4
July . . . . .	17.6	62	71	47	1	0	65 1900	Nil (e)	43 12/00	1.1
August . . . . .	20.6	66	73	53	2	0	84 1947	Nil (e)	80 2/47	0.8
September . . . . .	24.7	68	73	54	13	2	108 1942	Nil (e)	71 21/42	0.2
October . . . . .	27.7	68	72	60	50	5	339 1954	Nil (e)	95 28/56	0.0
November . . . . .	29.3	70	75	62	126	11	399 1938	10 1870	120 19/51	0.0
December . . . . .	30.5	75	83	65	243	16	616 1974	25 1934	279 25/74	0.0
Year { Totals . . . . .	..	..	..	..	1,536	97	..	..	..	2.5
Year { Averages . . . . .	25.9	71	..	..	..	..	..	..	..	..
Year { Extremes . . . . .	..	..	89	47	..	..	1013 3/77	Nil (f)	296 7/1/1897	..

(a) Records to 1966 at Aerodrome. (b) 1882 to 1938 at Post Office. (c) 1869 to 1962 at Post Office; 8 years missing. (d) Highest or lowest at either Post Office, Aerodrome or Regional Office Sites. (e) Various years. (f) April to October. Various years.

Figures such as 2/82, 26/42, etc., indicate in respect of the month of reference, the day and year of occurrence. Dates marked with an asterisk(\*) relate to nineteenth century.

## CLIMATIC DATA: ADELAIDE, SOUTH AUSTRALIA

(Lat. 34° 46' S., Long. 138° 35' E. Height above M.S.L. 43 metres)

## BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 22 metres)					Mean amt evapo- ration (mm)	No. days thun- der	Mean daily amt clouds 9 a.m., 3 p.m., 9 p.m., (a)	No. of clear days
		Average (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction					
					9 a.m.	3 p.m.				
No. of years of record .	119	20(b)	20(b)	59	30(c)	30(c)	9	104	108	61
January . . . . .	1,013.2	12.8	32.2	12/70	116	SW	261	1.5	3.0	12.0
February . . . . .	1,014.3	12.1	28.8	25/67	106	NE	224	1.1	3.0	10.7
March . . . . .	1,017.2	11.4	30.7	24/64	126	S	180	0.8	3.3	10.7
April . . . . .	1,019.8	11.4	37.4	10/56	130	NE	126	1.0	4.2	6.8
May . . . . .	1,020.1	11.3	37.8	19/53	113	NE	80	1.0	4.7	4.5
June . . . . .	1,019.8	11.6	29.7	16/70	108	NE	57	0.9	5.0	3.8
July . . . . .	1,019.9	11.8	32.9	13/64	148	NE	61	0.8	4.9	3.5
August . . . . .	1,019.0	12.8	38.2	8/55	121	NE	76	1.1	4.2	4.7
September . . . . .	1,017.6	13.2	34.9	16/65	111	NNE	113	1.3	4.3	5.5
October . . . . .	1,016.0	13.6	35.4	1/68	121	NNE	169	1.9	4.2	5.6
November . . . . .	1,015.1	13.9	36.3	14/68	130	SW	202	2.0	3.9	6.5
December . . . . .	1,013.3	13.5	31.1	18/69	121	SW	247	1.5	3.4	8.8
Year { Totals . . . . .	1,017.1	..	..	..	..	..	1,795	14.9	..	83.1
Averages . . . . .	1,017.1	..	..	..	NE	SW	..	..	4.0	..
Extremes . . . . .	..	..	38.2	8/65	148	..	..	..	..	..

(a) Scale 0-8. (b) Records of cup anemometer. (c) Standard 30 years normal (1931-1960).

## TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (°Celsius)			Extreme air temperature (°Celsius)		Extreme temperature (°Celsius)		Mean daily hours sunshine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record .	119	119	119	119	119	54(a)	115	94
January . . . . .	29.6	16.4	23.0	47.6	12/39	7.3	21/84*	9.9
February . . . . .	29.4	16.6	23.0	45.3	12/99*	7.5	23/18	9.3
March . . . . .	26.9	15.1	21.0	43.6	9/34	6.6	21/33	7.9
April . . . . .	22.7	12.7	17.7	37.0	5/38	4.2	15/59*	6.0
May . . . . .	18.7	10.3	14.5	31.9	4/21	2.7	(b)	4.8
June . . . . .	15.8	8.3	12.1	25.6	4/57	0.3	(c)	4.2
July . . . . .	15.0	7.3	11.1	26.6	29/75	0.0	24/08	4.3
August . . . . .	16.4	7.8	12.1	29.4	31/11	0.2	17/59*	5.3
September . . . . .	18.9	9.0	14.0	35.1	30/61	0.4	4/58*	6.2
October . . . . .	22.0	10.9	16.5	39.4	21/22	2.3	20/58*	7.2
November . . . . .	25.2	12.9	19.1	45.3	21/65*	4.9	2/09	8.6
December . . . . .	27.8	14.9	21.4	45.9	29/31	6.1	(d)	9.4
Year { Averages . . . . .	22.4	11.8	17.1	47.6	..	0.0	..	6.9
Extremes . . . . .	..	..	..	12/1/39	24/7/08	82.3	-6.1	..
						18/1/62	24/6/44	

(a) Discontinued 1934, incomplete 1931-1934. (b) 26/1895 and 24/04. (c) 27/1876 and 24/44. (d) 16/1861 and 4/06.

## HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)						Fog Mean No. days		
		Mean	Highest mean	Lowest mean	Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day				
No. of years of record .	108	108	108	108	137	137	137	137	137	137	137	76	
January . . . . .	11.9	41	59	29	20	4	84	1941	Nil	(a)	58	2/89*	0.0
February . . . . .	12.5	44	61	30	21	4	155	1925	Nil	(a)	141	7/25	0.0
March . . . . .	12.0	47	62	29	24	5	117	1878	Nil	(a)	89	5/78*	0.0
April . . . . .	11.5	57	72	37	44	9	154	1971	Nil	1945	80	5/60*	0.0
May . . . . .	10.8	67	77	49	69	13	197	1875	3	1934	70	1/53*	0.4
June . . . . .	10.0	75	84	63	72	15	218	1916	6	1958	54	1/20	1.1
July . . . . .	9.5	76	87	66	67	16	138	1890	10	1899	44	10/65*	1.3
August . . . . .	9.7	70	80	54	62	16	157	1852	8	1944	57	19/51*	0.6
September . . . . .	10.0	61	72	44	51	13	148	1923	7	1951	40	20/23	0.2
October . . . . .	10.2	52	67	29	44	11	133	1949	1	1969	57	16/08	0.0
November . . . . .	10.5	45	64	31	31	8	113	1839	1	1963	75	12/60	0.0
December . . . . .	11.3	42	56	31	26	6	101	1861	Nil	1904	61	23/13	0.0
Year { Totals . . . . .	..	..	..	..	531	120	..	..	..	..	..	..	3.6
Averages . . . . .	10.5	56	87	29	..	..	..	..	..	..	..	..	..
Extremes . . . . .	..	..	87	29	..	..	218	..	Nil	(b)	141	..	..
							6/1916				7/2/25		

(a) Various years. (b) December to April, various years.

Figures such as 3/55, 21/84, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk(\*) relate to nineteenth century.

## CLIMATIC DATA: BRISBANE, QUEENSLAND

(Lat. 27° 28' S., Long. 153° 2' E. Height above M.S.L. 41 metres)

## BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 32 metres)					Mean amt evapo- ration (mm)	No. days thun- der	Mean daily amt clouds	
		Average (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction				9 a.m., 3 p.m., 9 p.m. (a)	No. clear days
					9 a.m.	3 p.m.				
No. of years of record .	89	60	60	60	25(b)	25(b)	9	89	84	68
January . . . . .	1,011.7	12.1	31.8	23/47	SE	ENE	189	4.6	4.6	3.3
February . . . . .	1,012.5	11.9	37.3	21/54	SSW	ENE	150	3.7	4.8	2.4
March . . . . .	1,014.6	11.4	32.7	1/29	SSW	ESE	149	2.3	4.3	5.6
April . . . . .	1,017.3	10.5	26.8	3/25	SW	ESE	127	1.4	3.6	7.8
May . . . . .	1,018.3	9.8	28.8	17/26	SW	WSW	89	0.6	3.3	10.0
June . . . . .	1,018.5	10.0	30.5	14/28	SW	WSW	70	0.5	3.3	10.5
July . . . . .	1,018.8	9.7	35.4	13/54	SW	WSW	77	0.4	2.9	13.3
August . . . . .	1,018.8	10.0	23.8	4/35	SW	NE	105	1.4	2.6	13.5
September . . . . .	1,017.6	10.5	25.9	1/48	SW	NE	133	2.8	2.8	12.4
October . . . . .	1,015.9	11.1	25.3	1/41	SSW	NE	168	4.4	3.5	8.5
November . . . . .	1,014.1	11.4	24.9	10/28	SE	NE	191	5.7	3.9	6.1
December . . . . .	1,012.1	11.9	31.3	15/26	SSE	NE	209	6.6	4.3	4.5
Year { Totals . . . . .	1,015.9	10.8	..	..	..	..	1,656	34.2	..	97.7
Averages . . . . .	..	..	35.7	..	..	..	..	..	3.6	..
Extremes . . . . .	..	..	..	..	128	..	..	..	..	..
				21/2/54						

(a) Scale 0-8.

(b) 1950-1974.

## TEMPERATURE AND SUNSHINE

	Air temperature daily readings (°Celsius)			Extreme air temperature (°Celsius)		Extreme temperature (°Celsius)		Mean daily hours sun-shine
Month	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record .	89	89	89	89	89	50(a)	89	67
January . . . . .	29.4	20.6	25.0	43.2	14.9	76.2	9.9	7.5
February . . . . .	28.9	20.4	24.7	40.9	14.7	74.0	9.5	7.0
March . . . . .	27.8	19.2	23.5	38.8	11.3	72.5	7.4	6.8
April . . . . .	26.0	16.4	21.2	36.1	6.9	67.7	2.6	7.1
May . . . . .	23.1	13.1	18.0	32.4	4.8	63.9	-1.2	6.8
June . . . . .	20.8	10.7	15.7	31.6	2.4	57.8	-3.7	6.6
July . . . . .	20.3	9.4	14.9	29.1	2.3	63.4	-4.5	7.0
August . . . . .	21.8	10.0	15.9	32.8	2.7	61.1	-2.7	7.8
September . . . . .	24.0	12.7	18.3	38.3	4.8	68.6	-0.9	8.3
October . . . . .	26.1	15.8	20.9	40.7	6.3	69.7	1.6	8.2
November . . . . .	27.8	17.9	22.9	41.2	9.2	72.4	3.8	8.2
December . . . . .	29.1	19.6	24.5	41.1	13.5	74.4	9.5	8.1
Year { Averages . . . . .	25.4	15.5	20.5	43.2	2.3	76.2	-4.5	7.5
Extremes . . . . .	..	..	..	26/1/1940	..	2/1/1937	11/7/1890	..

(a) 1887-1926, 1936-March 1947.

(b) 12/1894 and 2/1896.

## HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)					Fog Mean No. days	
		Mean	Highest mean	Lowest mean	Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day		
No. of years of record .	64	89	89	89	124	116	123	123	123	89	
January . . . . .	21.7	65	79	53	167	13	872	1974	8 1919	465 21/87*	0.5
February . . . . .	22.0	69	82	55	161	14	1,026	1893	15 1849	270 6/31	0.6
March . . . . .	20.9	71	85	56	144	15	865	1870	Nil 1849	284 14/08	1.1
April . . . . .	17.5	70	80	56	88	11	388	1867	Nil 1944	178 3/72	2.1
May . . . . .	14.3	71	85	59	69	9	352	1876	Nil 1846	143 9/79*	3.0
June . . . . .	12.1	72	84	54	69	8	647	1967	Nil 1847	283 12/67	2.9
July . . . . .	11.1	70	88	53	54	7	330	1973	Nil (c)	193 20/65	3.0
August . . . . .	11.7	66	80	53	48	7	373	1879	Nil (b)	124 12/87*	3.6
September . . . . .	13.8	63	76	47	48	8	138	1886	3 1907	79 12/65	2.5
October . . . . .	16.0	60	72	48	74	9	456	1972	(c) 1948	136 25/49	1.2
November . . . . .	18.1	59	72	45	95	10	315	1917	Nil 1842	143 8/66*	0.5
December . . . . .	20.1	61	70	51	129	12	441	1942	9 1865	168 28/71*	0.3
Year { Totals . . . . .	..	..	..	..	1,157	123	..	..	..	..	21.3
Averages . . . . .	16.6	66	..	..	..	..	..	..	..	..	..
Extremes . . . . .	..	..	88	45	..	..	1,026	..	Nil	465	..
							2/1893	Various		21/1/1887	

(a) 1841 and 1951.

(b) 1862, 1869 and 1880.

(c) Less than 1 mm.

Figures such as 23/47, 4/93, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk (\*) relate to nineteenth century.

## CLIMATIC DATA: SYDNEY, NEW SOUTH WALES

(Lat. 33° 52' S., Long. 151° 12' E. Height above M.S.L. 42 metres)

## BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 22 metres)						Mean amt evapora- (mm)	No. days thun- der	Mean daily amt clouds 9 a.m., 3 p.m., 9 p.m. (a)	No. clear days
		Average (km/h)	Highest mean speed in one day (km/h)	High- est gust speed (km/h)	Prevailing direction						
					9 a.m.	3 p.m.					
No. of years of record .	66	25(b)	25(b)	25(b)	25(b)	25(b)	84(c)	56	114	65	
January . . . . .	1,012.7	12.3	30.3	10/49	150	NE	NE	189	3.3	4.7	4.9
February . . . . .	1,014.2	11.6	30.3	18/57	101	NE	ENE	142	2.5	4.8	4.5
March . . . . .	1,016.4	10.5	33.3	10/44	93	WNW	ENE	142	1.7	4.4	5.7
April . . . . .	1,018.3	10.2	36.2	24/44	116	W	ENE	108	1.3	4.1	7.3
May . . . . .	1,018.7	10.5	33.8	18/55	101	W	ENE	85	0.9	3.9	7.7
June . . . . .	1,018.9	11.6	36.0	10/47	135	W	WSW	58	0.8	4.0	6.0
July . . . . .	1,018.5	11.5	34.3	20/51	106	W	WSW	73	0.8	3.5	10.5
August . . . . .	1,017.9	12.1	39.6	9/51	109	WNW	WNW	112	1.4	3.3	10.4
September . . . . .	1,017.0	11.6	35.1	23/42	113	WNW	NE	150	1.8	3.5	9.1
October . . . . .	1,015.1	12.3	39.4	1/57	153	WNW	ENE	203	2.7	4.1	6.5
November . . . . .	1,013.4	12.4	31.9	21/54	114	WNW	ENE	190	3.6	4.5	5.2
December . . . . .	1,012.1	12.3	36.2	11/52	121	NE	ENE	225	3.8	4.6	4.8
Year { Totals . . . . .	1,016.1	11.6	..	..	..	..	..	1,677	24.7	..	84.7
Year { Averages . . . . .	..	..	..	..	..	WNW	ENE	..	..	4.2	..
Year { Extremes . . . . .	..	..	39.6	153	..	..	..	..	..	..	..
			9/8/51								

(a) Scale 0-8.

(b) Years 1938-1962 inclusive.

(c) Richmond records.

## TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (°Celsius)			Extreme air temperature (°Celsius)		Extreme temperature (°Celsius)		Mean daily hours sun-shine				
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass					
No. of years of record .	117	117	117	117	117	84(a)	117	55				
January . . . . .	25.7	18.3	22.0	45.3	14/39	10.6	18/49	73.5	26/15	6.5	6/25	7.2
February . . . . .	25.4	18.4	21.9	42.1	8/26	9.6	28/63*	76.3	14/39	6.0	22/33	6.8
March . . . . .	24.5	17.3	20.9	39.2	3/69*	9.3	14/86*	70.2	10/26	4.4	17/13	6.3
April . . . . .	22.1	14.5	18.3	33.0	(b)	7.0	27/64*	62.3	10/77*	0.7	24/09	6.2
May . . . . .	19.2	11.2	15.2	30.0	1/19	4.4	30/62*	54.3	1/96*	-1.5	25/17	5.8
June . . . . .	16.6	9.1	12.8	26.9	11/31	2.1	22/32	52.1	2/23	-2.2	22/32	5.2
July . . . . .	15.8	7.8	11.8	25.7	22/26	2.2	12/90*	51.9	19/77*	-4.4	4/93*	6.2
August . . . . .	17.4	8.7	13.1	30.4	24/54	2.7	3/72*	65.0	30/78*	-3.3	4/09	6.8
September . . . . .	19.6	10.8	15.2	34.6	26/65	4.9	2/45	61.2	12/78*	-1.1	17/05	7.1
October . . . . .	21.9	13.3	17.6	37.4	4/42	5.7	6/27	66.8	20/33	0.4	9/05	7.3
November . . . . .	23.5	16.3	19.4	40.3	6/46	7.7	1/05	70.3	28/99*	1.9	21/67	7.6
December . . . . .	24.9	17.2	21.1	42.2	20/57	9.1	3/24	73.5	27/89*	5.2	3/24	7.4
Year { Averages . . . . .	21.4	13.6	17.4	..	..	..	..	..	..	..	..	6.7
Year { Extremes . . . . .	..	..	..	45.3	2.1	76.3	-4.4	..	..	..	..	..
				14/1/39		22/6/32		14/2/39		4/7/1893		

(a) Records discontinued 1946.

(b) 1/36 and 10/69.

## HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)					Fog Mean No. days			
		Mean	Highest mean	Lowest mean	Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day				
No. of years of record .	100	100	100	100	117	117	117	117	117	55			
January . . . . .	18.8	68	78	58	100	13	388	1911	6	1932	180	13/11	0.3
February . . . . .	19.2	70	81	60	115	13	564	1950	3	1939	226	25/73*	0.6
March . . . . .	18.3	74	85	62	131	14	521	1942	8	1965	281	28/42	1.5
April . . . . .	15.0	74	87	63	126	13	622	1861	2	1868	191	29/60*	2.1
May . . . . .	11.9	75	90	63	123	13	585	1919	4	1957	212	28/89*	3.1
June . . . . .	10.2	76	89	63	133	12	643	1950	4	1962	131	16/84*	2.7
July . . . . .	9.6	74	88	59	104	11	336	1950	2	1970	198	7/31	2.1
August . . . . .	9.5	68	84	54	81	11	378	1899	1	1885	140	22/71	1.7
September . . . . .	11.3	66	79	49	69	11	357	1879	2	1882	145	10/79*	0.9
October . . . . .	13.0	62	77	46	76	12	283	(a)	2	1971	162	13/02	0.6
November . . . . .	15.0	62	79	42	78	12	577	1961	2	1915	133	27/55	0.5
December . . . . .	17.6	64	77	51	79	13	402	1920	6	1913	121	13/10	0.4
Year { Totals . . . . .	..	..	..	..	1,215	148	..	..	..	..	..	..	16.3
Year { Averages . . . . .	14.1	69	..	..	..	..	643	6/1950	..	..	..	..	..
Year { Extremes . . . . .	..	..	90	42	..	..	..	..	1	8/1885	281	..	..
											28/3/1942		

(a) 1916 and 1959.

Figures such as 10/49, 28/63, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk(\*) relate to nineteenth century.

## CLIMATIC DATA: CANBERRA, AUSTRALIAN CAPITAL TERRITORY

(Lat. 35° 19' S., Long. 149° 11' E. Height above M.S.L. 571 metres)

## BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 10 metres)				Mean amt evaporation (mm)	No. days thunder	Mean daily amt clouds	
		Average (km/h)	Highest mean speed in one day (km/h)	Highest gust speed (km/h)	Prevailing direction			9 a.m., 3 p.m., 9 p.m. (a)	No. clear days
No. of years of record .	36	44(b)	44(b)	36(c)	36(c)	36(c)	8	36	36
January . . . . .	1,012.1	6.6	24 24/33	121	NW	NW	242	3.3	4.1
February . . . . .	1,013.1	6.1	25 24/33	104	NW	NW	194	3.0	4.4
March . . . . .	1,016.0	5.3	29 28/42	111	SE	NW	165	1.7	4.2
April . . . . .	1,018.8	5.0	30 8/45	106	NW	NW	109	0.8	4.2
May . . . . .	1,019.0	4.4	21 27/58	104	NW	NW	71	0.4	4.5
June . . . . .	1,021.0	4.8	26 2/30	96	NW	NW	46	0.2	4.6
July . . . . .	1,020.2	5.0	38 7/31	102	NW	NW	54	0.1	4.4
August . . . . .	1,018.5	5.9	25 25/36	113	NW	NW	77	0.8	4.4
September . . . . .	1,017.4	6.0	28 28/34	107	NW	NW	115	1.1	4.1
October . . . . .	1,014.8	6.5	23 12/57	119	NW	NW	165	2.2	4.4
November . . . . .	1,011.9	6.9	28 28/42	128	NW	NW	200	3.3	4.4
December . . . . .	1,010.7	6.9	26 11/38	106	NW	NW	259	3.4	4.1
Year { Totals . . . . .	1,016.1	5.8	..	..	NW	NW	1,697	20.3	..
Averages . . . . .	..	..	..	..	..	..	..	..	..
Extremes . . . . .	..	..	38 7/7/31	128	..	..	..	4.3	6.9

(a) Scale 0-8. (b) Recorded at Forestry and Timber Bureau, Yarralumla, where a cup anemometer is installed. (c) Recorded at Meteorological office, R.A.A.F. Fairbairn, where a Dines Pressure Tube anemometer is installed. (d) 1940-75. Formerly assessed over 37-year period at Yarralumla.

## TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings (°Celsius)			Extreme air temperature (°Celsius)		Extreme temperature (°Celsius)		Mean daily hours sunshine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record .	36	36	36	36	36	..	..	37
January . . . . .	27.5	12.9	20.2	41.4 31/68	1.8 1/56	..	-0.4 1/56	8.9
February . . . . .	26.6	12.6	19.6	42.2 1/68	3.0 16/62	..	0.2 17/70	8.2
March . . . . .	24.3	10.4	17.3	36.4 9/40	-1.1 24/67	..	-4.0 (a)	7.5
April . . . . .	19.6	6.5	13.1	32.6 12/68	-3.3 26/72	..	-8.3 24/69	6.9
May . . . . .	14.9	2.8	8.9	24.5 10/67	-7.5 30/76	..	-10.4 26/69	5.6
June . . . . .	12.0	0.8	6.4	20.1 3/57	-8.5 8/57	..	-13.4 25/71	4.8
July . . . . .	11.1	-0.3	5.4	19.7 29/75	-10.0 11/71	..	-15.1 11/71	5.1
August . . . . .	12.6	0.8	6.7	21.7 24/54	-7.8 6/74	..	-12.8 11/69	6.1
September . . . . .	15.8	2.7	9.3	28.6 26/65	-5.6 5/40	..	-10.6 12/71	7.4
October . . . . .	19.0	5.8	12.4	32.7 13/46	-3.3 4/57	..	-6.2 4/57	7.9
November . . . . .	22.2	8.2	15.1	38.8 19/44	-1.8 28/67	..	-6.3 28/67	8.7
December . . . . .	26.0	11.1	18.6	38.8 21/53	1.1 18/64	..	-3.9 18/64	9.1
Year { Averages . . . . .	19.3	6.2	12.7	..	..	..	..	7.2
Extremes . . . . .	..	..	..	42.2 1/2/68	-10.0 11/7/71	..	-15.1 11/7/71	..

(a) 30/58 and 24/67.

## HUMIDITY, RAINFALL, AND FOG

Month	Vapour pres- sure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rain/fall (millimetres)				Fog Mean No. days	
		Mean	Highest mean	Lowest mean	Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day		
No. of years of record .	36(a)	36	36	36	36	36	36	36	36	
January . . . . .	13.1	60	75	42	61	8	164 1941	1 1947	95 12/45	1.1
February . . . . .	14.0	65	81	53	59	7	145 1948	Nil 1968	69 20/74	1.2
March . . . . .	13.1	69	81	53	51	7	312 1950	1 1954	66 5/59	2.8
April . . . . .	10.7	75	84	38	50	8	164 1974	2 1942	75 2/59	4.1
May . . . . .	8.7	84	96	73	51	9	150 1953	1 1976	96 3/48	7.5
June . . . . .	7.1	85	97	73	39	9	126 1956	5 1971	45 25/56	7.6
July . . . . .	6.6	84	93	68	38	10	103 1960	4 1970	35 10/57	7.7
August . . . . .	7.1	80	92	58	47	12	161 1976	7 1944	48 29/74	5.0
September . . . . .	8.1	74	82	55	50	10	116 1970	6 1946	41 16/62	4.1
October . . . . .	10.0	67	82	50	73	12	148 1959	6 1940	105 21/59	3.1
November . . . . .	10.7	59	76	38	64	10	135 1961	13 1940	64 9/50	1.4
December . . . . .	12.3	59	74	43	56	8	215 1947	Nil 1967	87 30/48	0.6
Year { Totals . . . . .	..	..	..	..	639	110	..	..	..	46.2
Averages . . . . .	9.3	72	..	..	..	..	..	..	..	..
Extremes . . . . .	..	..	97	38	..	..	312 3/50	Nil (b)	105 21/10/59	..

(a) Formerly assessed over 38-year period at Forestry and Timber Bureau, Yarralumla. (b) 12/67 and 2/68.

Data shown in the above tables relate to the Meteorological Office, R.A.A.F., Fairbairn, except where otherwise indicated, and generally cover years up to 1976.

Figures such as 24/33, 31/68, etc., indicate, in respect of the month of reference, the day and year of the occurrence.

## CLIMATIC DATA: MELBOURNE, VICTORIA

(Lat. 37° 49' S., Long. 144° 58' E. Height above M.S.L. 35 metres)

## BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 28 metres)				Mean amt evaporation (mm)	No. days thunder	Mean daily amt clouds	
		Average (km/h)	Highest mean speed in one day (km/h)	Highest gust speed (km/h)	Prevailing direction			9 a.m., 9 p.m. (a)	No. clear days
No. of years of record .	119	36(b)	63	66	57	57	9	68	119
January . . . . .	1,012.8	12.8	34.0 27/41	106	S	S	228	1.7	4.1
February . . . . .	1,014.3	12.4	30.6 13/47	119	S	S	198	1.9	4.0
March . . . . .	1,016.8	11.3	29.0 3/61	106	N	S	155	1.3	4.3
April . . . . .	1,018.9	10.9	33.7 27/71	108	N	S	97	0.7	4.7
May . . . . .	1,019.1	11.4	33.0 4/61	116	N	N	59	0.4	5.2
June . . . . .	1,019.0	11.4	36.7 16/47	103	N	N	38	0.2	5.3
July . . . . .	1,018.6	12.8	36.9 24/70	109	N	N	47	0.2	5.2
August . . . . .	1,017.5	12.5	34.3 20/42	108	N	N	60	0.6	5.0
September . . . . .	1,016.0	12.7	34.0 15/64	111	N	S	91	0.8	4.8
October . . . . .	1,014.7	12.8	30.4 6/68	111	N	S	130	1.6	4.8
November . . . . .	1,013.9	13.3	35.8 8/71	114	SW	S	161	1.9	4.9
December . . . . .	1,012.4	13.1	33.8 12/52	100	S	S	209	2.2	4.5
Year { Totals . . . . .	1,016.2	12.3	.. ..	..	..	1,468	13.4	..	48.0
Averages . . . . .	..	..	.. ..	..	..	..	..	..	..
Extremes . . . . .	..	..	36.9	119	..	..	..	..	..
24/7/70									

(a) Scale 0-8. (b) Early records not comparable. (c) Records to 1966.

## TEMPERATURE AND SUNSHINE

	Air temperature daily readings (°Celsius)			Extreme air temperature (°Celsius)		Extreme temperature (°Celsius)		Mean daily hours sun-shine
Month	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record .	120	120	120	120	120	86(a)	116	52(b)
January . . . . .	25.8	13.9	19.9	45.6 13/39	5.6 28/85*	81.4 14/62*	-1.0 28/85*	8.1
February . . . . .	25.7	14.2	19.9	43.1 7/01	4.6 24/24	75.3 15/70*	-0.6 6/91*	7.5
March . . . . .	23.7	12.8	18.3	41.7 11/40	2.8 17/84*	73.6 1/68*	-1.7 (c)	6.6
April . . . . .	20.1	10.5	15.3	34.9 5/38	1.6 24/88*	66.7 8/61*	-3.9 23/97*	5.1
May . . . . .	16.5	8.3	12.4	28.7 7/05	-1.2 29/16	61.4 2/59*	-6.1 26/16	3.9
June . . . . .	13.9	6.6	10.3	22.4 2/57	-2.2 11/66	53.9 11/61*	-6.7 30/29	3.4
July . . . . .	13.3	5.7	9.5	23.1 30/75*	-2.8 21/69*	52.1 27/80*	-6.4 12/03	3.7
August . . . . .	14.8	6.4	10.6	25.0 20/85*	-2.1 11/63*	58.6 29/69*	-5.9 14/02	4.6
September . . . . .	17.1	7.6	12.4	31.4 28/28	-0.6 3/40	61.2 20/67*	-5.1 8/18	5.5
October . . . . .	19.5	9.2	14.4	36.9 24/14	0.1 3/71*	67.9 28/68*	-4.0 22/18	5.9
November . . . . .	21.8	10.8	16.3	40.9 27/94*	2.4 2/96*	70.9 29/65*	-4.1 2/96*	6.5
December . . . . .	24.1	12.6	18.4	43.7 15/76	4.4 4/70*	76.8 20/69*	0.7 1/04	7.3
Year { Averages . . . . .	19.9	9.9	14.8	45.6 ..	-2.8 ..	81.4 ..	-6.7 ..	5.7
Extremes . . . . .	..	..	..	13/1/39	21/7/69	14/1/62	30/6/29	..

(a) Discontinued 1946. (b) Discontinued 1967. (c) 17/1884 and 20/1897.

## HUMIDITY, RAINFALL, AND FOG

Month	Vapour pressure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)				Fog Mean No. days
		Mean	Highest mean	Lowest mean	Mean mthly	No. of days of rain	Greatest monthly	Least monthly	
No. of years of record .	68	68	68	68	120	120	120	120	118
January . . . . .	13.1	61	68	50	48	8	176 1963	(a) 1932	108 29/63
February . . . . .	14.1	63	77	48	50	7	238 1972	(a) 1965	87 26/46
March . . . . .	13.3	66	79	50	54	9	191 1911	4 1934	90 5/19
April . . . . .	11.7	72	82	66	59	11	195 1960	Nil 1923	80 23/66
May . . . . .	10.3	79	88	69	57	14	142 1942	4 1934	51 15/74
June . . . . .	9.3	83	92	73	50	14	114 1859	8 1858	44 22/04
July . . . . .	8.9	81	87	73	49	15	178 1891	15 1902	74 12/91*
August . . . . .	9.1	75	82	65	50	15	111 1939	12 1903	54 17/81*
September . . . . .	9.5	68	76	60	59	14	201 1916	13 1907	59 23/16
October . . . . .	10.5	63	72	52	68	14	193 1869	7 1914	61 21/53
November . . . . .	11.3	61	70	52	59	12	206 1954	6 1895	73 21/54
December . . . . .	12.5	60	69	48	58	10	182 1863	1 1972	100 4/54
Year { Totals . . . . .	11.1	..	..	..	661	143	.. ..	.. ..	19.3
Averages . . . . .	..	69	..	..	..	..	238 2/72	Nil 4/23	108 ..
Extremes . . . . .	..	..	92	43	..	..	..	..	..
29/1/63									

(a) Less than 1 mm.

Figures such as 27/41, 28/85, etc., indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk(\*) relate to nineteenth century.

## CLIMATIC DATA: HOBART, TASMANIA

(Lat. 42° 53' S., Long. 147° 20' E. Height above M.S.L. 54 metres)

## BAROMETER, WIND, EVAPORATION, THUNDER, CLOUDS, AND CLEAR DAYS

Month	Mean of 9 a.m. and 3 p.m. atmospheric pressure reduced to mean sea level (mb)	Wind (height of anemometer 12 metres)						Mean amt evaporation (mm)	No. days thunder	Mean daily amt clouds	
		Average (km/h)	Highest mean speed in one day (km/h)	Highest gust speed (km/h)	Prevailing direction		9 a.m., 3 p.m.			No. clear days	
					9 a.m.	3 p.m.					
No. of years of record .	90	63	64	84	30(b)	30(b)	10(c)	64	90	30(b)	
January . . . . .	1,010.6	12.6	33.5	30/16	130	NNW	SSE	167	1.0	5.0	1.9
February . . . . .	1,012.9	11.5	40.6	4/27	121	NNW	SSE	135	1.0	4.9	2.3
March . . . . .	1,014.3	11.0	34.4	13/38	127	NW	SSE	109	0.7	4.8	2.4
April . . . . .	1,015.5	10.9	38.8	9/52	121	NW	W	70	0.3	5.0	1.7
May . . . . .	1,015.4	10.4	35.4	21/65	135	NNW	NW	38	0.0	5.0	2.4
June . . . . .	1,015.2	10.2	38.2	27/20	132	NW	NW	22	0.0	5.0	2.4
July . . . . .	1,014.0	10.7	36.9	22/53	129	NNW	NNW	26	0.0	4.8	2.0
August . . . . .	1,012.8	10.9	41.0	19/26	140	NNW	NW	44	0.1	4.8	2.1
September . . . . .	1,011.4	12.5	43.1	28/65	150	NNW	NW	73	0.1	4.9	1.5
October . . . . .	1,010.3	12.6	32.4	3/65	140	NNW	SW	107	0.4	5.2	1.0
November . . . . .	1,009.8	12.8	34.1	18/15	135	NNW	S	123	0.6	5.3	1.3
December . . . . .	1,009.4	12.4	37.7	1/34	122	NNW	SSE	150	0.8	5.3	1.1
Year { Totals . . . . .								1,064	5.1		22.1
Averages . . . . .	1,012.6	11.5				NNW	W			5.0	
Extremes . . . . .			43.1	28/9/65	150						

(a) Scale 0-8. (b) Standard thirty years normal (1911-1940). (c) Class "A" American pan.

## TEMPERATURE AND SUNSHINE

Month	Air temperature daily readings* (°Celsius)			Extreme air temperature (°Celsius)		Extreme temperature (°Celsius)		Mean daily hours sun shine
	Mean max.	Mean min.	Mean	Highest	Lowest	Highest in sun	Lowest on grass	
No. of years of record .	92	92	92	92	92	57(a)	87	79
January . . . . .	21.4	11.5	16.5	40.8	4.5	71.1	19/97*	7.9
February . . . . .	21.5	11.8	16.7	40.2	12/99*	73.9	24/98*	7.0
March . . . . .	20.0	10.6	15.3	37.3	13/40	66.1	26/44	6.4
April . . . . .	17.1	8.7	12.9	30.6	1/41	61.1	18/93*	5.0
May . . . . .	14.2	6.7	10.5	25.5	5/21	53.3	(d)	4.3
June . . . . .	11.8	5.1	8.5	20.6	1/07	50.0	12/94*	3.9
July . . . . .	11.4	4.4	7.9	21.0	30/75	49.4	12/93*	4.3
August . . . . .	12.8	5.0	8.9	22.0	28/14	54.4	—/87*	5.0
September . . . . .	14.9	6.2	10.6	28.2	29/73	58.9	23/93*	5.9
October . . . . .	16.7	7.5	12.1	33.4	24/14	68.9	9/93*	6.3
November . . . . .	18.5	9.0	13.8	36.8	26/37	55.6	19/92*	7.0
December . . . . .	20.2	10.5	15.4	40.7	30/97*	71.9	10/39	7.2
Year { Averages . . . . .	16.7	8.1	12.4	..	..	73.9	..	5.9
Extremes . . . . .	..	..	..	40.8	—2.8	73.9	—7.7	..
				4/1/1976	25/6/72	24/2/1968	24/6/1963	

(a) Period 1934-1938 not comparable; records discontinued 1946. (b) 09/1937 and 11/1937. (c) 05/1886 and 13/1905. (d) —/1899 and —/1893. (e) 1/1886 and 1/1899. (f) 11/1895 and 7/1973.

## HUMIDITY, RAINFALL, AND FOG

Month	Vapour pres- sure mean 9 a.m. (mb)	Rel. hum. (%) at 9 a.m.			Rainfall (millimetres)					Fog Mean No. days	
		Mean	Highest mean	Lowest mean	Mean mthly	Mean No. of days of rain	Greatest monthly	Least monthly	Greatest in one day		
No. of years of record .	77(a)	81	81	81	93	93	92	92	131	64	
January . . . . .	11.0	58	81	45	49	11	150	1893	4 1958	75 30/16	0.3
February . . . . .	11.7	62	83	49	42	10	171	1964	3 1914	56 1/54	0.1
March . . . . .	11.0	65	78	52	47	11	255	1946	7 1943	88 17/46	0.3
April . . . . .	10.0	70	84	57	55	12	248	1960	2 1904	133 23/60	0.3
May . . . . .	8.8	75	86	61	49	14	214	1958	4 1913	47 3/73	1.1
June . . . . .	7.9	78	91	61	59	14	238	1954	7 (c)	147 7/54	1.7
July . . . . .	7.6	78	87	72	54	15	157	1974	4 1950	64 18/22	1.4
August . . . . .	7.9	73	86	59	49	16	161	1946	8 1892	58 14/90*	0.7
September . . . . .	8.3	66	81	52	52	15	201	1957	10 1951	156 15/57	0.2
October . . . . .	9.1	62	74	52	64	17	193	1947	10 1914	66 4/06	0.1
November . . . . .	9.6	59	73	49	56	14	188	1885	9 (d)	94 30/85*	0.1
December . . . . .	10.6	58	73	42	57	13	196	(b)	5 (e)	85 5/41	0.1
Year { Totals . . . . .	..	..	..	..	633	162	..	..	..	..	6.1
Averages . . . . .	9.5	67	..	..	..	..	..	..	..	..	..
Extremes . . . . .	..	..	91	42	..	..	255	..	2	156	..
							3/1946		4/1904	15/9/57	

(a) 1894-1970. (b) 1897 and 1916. (c) 1886 and 1967. (d) 1919 and 1921. (e) 1897, 1915 and 1931.

Figures such as 30/16, 12/99, etc. indicate, in respect of the month of reference, the day and year of the occurrence. Dates marked with an asterisk(\*) relate to nineteenth century.

## REFERENCES

Bureau of Meteorology (Melbourne), Department of Science.

- Ashton, H. T. . . . 1964 *Meteorological Data for Air Conditioning in Australia.* Bulletin 47.
- Foley, J. C. . . . 1945 (i) *Frost in the Australian Region.* Bulletin 32  
1957 (ii) *Droughts in Australia.* Bulletin 43.
- Gaffney, D. O. . . . 1971 (i) *Snowy Mountains Median Annual Rainfall* Working Paper 139  
1971 (ii) *Seasonal Rainfall Zones in Australia* Working Paper 141
- Gibbs, W. J. and Maher, J. V. 1967 *Rainfall Deciles as Drought Indicators.* Bulletin 48.
- Hounam, C. E. . . . 1969 *Climate and Air Conditioning Requirements in Sparsely Occupied Areas of Australia.* Working Paper 30/78.
- Whittingham, H. E. . . 1964 *Extreme Wind Gusts in Australia.* Bulletin 46.

## Other Publications

- American Society of Heating, 1960 *Physiological Principles. In Heating, Ventilating and Air Refrigerating and Air Conditioning Engineers* Vol. 38, (Washington; the Society).
- Gaffney, D. O. . . . 1970 *Rainfall*  
1973 *Temperatures*  
1973 *Climate*  
Commentaries prepared for *Atlas of Australian Resources*, Second Series, Department of Minerals and Energy, Canberra.
- 1975 *Rainfall Deficiency and Evaporation in relation to Drought in Australia.* ANZAAS Congress, Canberra.
- Lee, D. H. K. and Henschel, A. 1963 *Evaluation of Environment in Shelters.* U.S. Department of Health, Education and Welfare (Cincinnati: the Department).