


ABS DATA INTEGRATION & DIGITAL SERVICES

ABS DATA INTEGRATION & DIGITAL SERVICES

DataLab

DataLab Safe Researcher Training

Part 3: Safe Outputs and statistical disclosure control



ABS DATA INTEGRATION & DIGITAL SERVICE

Training Outcomes



Understand concepts in statistical disclosure control

 Know how to prepare safe outputs that are nondisclosive

Statistical Disclosure Control



- What is SDC?
 - Checking for disclosure risk in results leaving the 'safe settings'
 - Applying treatments where disclosure risk is too high
- Principles of SDC
 - Precautionary
 - Balancing risk and utility
 - Consistent with good research
- SDC in practice
 - Output rules in the User Guide



Why are safe outputs so important





Legal Only release data that is 'not likely to identify' **Ongoing Data Sharing** Data Custodians have confidence the sharing data won't lead to disclosure

Ongoing Data Collection People and businesses have confidence their information is handled appropriately **Risk management** Only the results that need to be are removed from the secure environment

Outputs from the DataLab



Everything that leaves the DataLab must first be checked by the ABS DataLab clearance team



Producing safe outputs

- Follow the DataLab output rules
 - Provide evidence
 - Apply treatments
- Principles-based approach to less common analysis
- Requesting exceptions to the standard rules
 - These will be escalated expect delays
 - You will need to show evidence that it's important, non-disclosive, and uncommon
 - Any exceptions are non-precedent setting

Main output rules





Rule of 10
 Dominance
 Model-specific rules

4. Quantiles5. Group Disclosure6. Secondary Contributors

Output treatment options



- Treatment should change the output to the point at which is passes the rules
 - Combine categories in tables
 - Round cells to the nearest 5, 10, 100, 1000, 10000, ...
 - Perturb/add noise to each cell
 - Use words to describe the output *"The relative proportions for population X is similar to population Y."*
 - Suppress problematic cells (remember secondary)





WHY? To prevent the re-identification of units in cells with small counts

WHERE? Rule applies to most outputs (table cells, sums/means, counts used to create charts etc)

Counts of less than 10 should also not be able to be derived from the available data

Each cell should have at least **10 contributing** units

Example 1 – Rule of 10



Table: Fortnightly income for persons living on Norfolk Island aged 20-24Source: Census 2021

	Count	%
Nil income	10	5.6
\$1-\$500	8	4.5
\$501-\$1000	40	22.5
\$1001-\$1500	40	22.5
\$1501-\$2000	45	25.3
\$2001-\$2500	25	14.0
\$2501 or more	10	5.6
Total	178	100.0

Example 1 – Rule of 10 TREATED



Table: Fortnightly income for persons living on Norfolk Island aged 20-24Source: Census 2021

	Count	%
Nil income	10	5.6
\$1-\$500	n/a	n/a
\$501-\$1000	40	22.5
\$1001-\$1500	40	22.5
\$1501-\$2000	45	25.3
\$2001-\$2500	25	14.0
\$2501 or more	n/a	n/a
Total	178	100.0

	Count	%
Nil income - \$500	18	10.1
\$501-\$1000	40	22.5
\$1001-\$1500	40	22.5
\$1501-\$2000	45	25.3
\$2001-\$2500	25	14.0
\$2501 or more	10	5.6
Total	178	100.0

Example 2 – Rule of 10



Average Weekly coffees by age group – Persons studying at University

Table 1 – Age groups as per the US Standard

Table 2 – Age groups as per the Australian Standard

		Age Group				Age Group	
Coffees per week	<21	21 and over	Total	Coffees per week	<18	18 and over	Total
0	135	124	259	0	120	139	259
1-2	132	99	231	1-2	126	105	231
3-5	99	92	191	3-5	85	106	191
6-9	100	138	238	6-9	76	162	238
10 or more	91	120	211	10 or more	76	135	211
Not stated	127	79	206	Not stated	117	89	206

Example 2 – Rule of 10 TREATED



Average Weekly coffees by age group – Persons studying at University

Table 1 – Age groups as per the US Standard

	Age Group			
Coffees per week	<21	21 and over	Total	
0	140	120	260	
1-2	130	100	230	
3-5	100	90	190	
6-9	100	140	240	
10 or more	100	120	210	
Not stated	130	80	210	

Table 2 – Age groups as per the Australian Standard

	Age Group			
Coffees per week	<18	18 and over	Total	
0	120	140	260	
1-2	130	110	230	
3-5	90	110	190	
6-9	90	160	240	
10 or more	80	140	210	
Not stated	120	90	210	





WHY? To prevent the re-identification of units that contribute a large percentage of a cell's total value

WHERE? Applies mainly to sums/totals and means

The **largest** contributor must contribute less than 50% The **two largest** contributors must contribute less than 67%

Example 3 - Dominance



Total turnover (\$M) of all pharmacies by Local Government Area

LGA Code	Total Turnover	No. of Businesses
1	1.65	12
2	0.94	11
3	3.22	20
4	2.10	10
5	2.05	16
Total	9.96	69

Example 3 - Dominance





Total turnover (\$M) of all pharmacies by Local Government Area

LGA Code	Total Turnover	No. of Businesses	Turnover of largest business	Turnover of 2 nd largest business	Proportion from largest business to total	Proportion from largest two businesses to total
1	1.65	12	0.66	0.59	40%	76%
2	0.94	11	0.14	0.13	15%	29%
3	3.22	20	1.77	0.32	55%	65%
4	2.10	10	0.74	0.46	35%	57%
5	2.05	16	0.86	0.29	42%	56%
Total	9.96	69	1.79	0.80	18%	26%

ABS DATA INTEGRATION & DIGITAL SERVICES

Example 3 – Dominance - TREATED





Total turnover (\$M) of all pharmacies by Local Government Area

LGA Code	Total Turnover	No. of Businesses		Turnover of 2 nd largest business	Proportion from largest business	Proportion from largest two businesses
1&3	4.87	32	1.77	0.66	36%	50%
2	0.94	11	0.14	0.13	15%	29%
4	2.1	10	0.74	0.46	35%	57%
5	2.05	16	0.86	0.29	42%	56%
Total	9.96	69	1.79	0.8	18%	26%
			OR			

"Total turnover ranking for the five LGAs of interest were (from largest to smallest): LGA 3, 4, 5, 1 and then 2."

Model-specific rules



WHY? Designed to prevent the re-identification of units using overfitted models and/or residuals

WHERE? All modelling outputs

The model should have at least **10 degrees of freedom** The **R-squared** for least squares regression should be <= 0.9 **Individual residuals** cannot leave the DataLab Extra rules when the independent variables are all categorical (contact the ABS)

Example 4 – Model-specific rules



Linear regression that looks at personal income as a function of a range of variables.

Variable	Model 1	Model 2	Model 3	Model 4
Sex	11.34	8.35	8.12	8.33
Age	1.61	1.56	1.55	1.55
SEIFA (index value)		17.28	17.33	17.33
Completed Yr 12			-6.76	-7.93
Has Bachelor Degree				2.36
Constant	36.85	-9.88	-5.23	-5.27
Ν	371	371	371	371
r ²	0.23	0.78	0.79	0.97

Minimum contributors for quantiles



WHY? To prevent the re-identification of units in from a group with small counts

WHERE? Any quantiles, maximum, minimum, range

Each "bin" must have **at least 5 contributors** No **minimums** or **maximums** out of DataLab

	Minimum contributors
Percentiles	500
Deciles	50
Quartiles	20
Median	10

Example 5 - Quantiles





Age	Count
0	11
1	0
2	4
3	6
4	14
5	17
6	11
7	17
8	9
9	6
10	2
11	1
12	0
13	0
14	2
Total	100

	Original	Requirement	Treated
Minimum	0	Min 10 in cell	ОК - 0
5 th percentile	0	100 total contributors	ОК - 0
Median	5	10 total contributors	ОК - 5
95 th percentile	9.5	100 total contributors	ОК — 9.5
99 th percentile	14	500 total contributors	Cannot clear
Maximum	14	Min 10 in cell	Cannot clear

ABS DATA INTEGRATION & DIGITAL SERVICES

Group Disclosure Rule



WHY? To protect the disclosure of a previously unknown attribute of an individual or business from a given group, where that group has a common feature

WHERE? Totals, means, proportions, counts

Particularly important where there is a risk of adverse consequences to the group

No cells should contain more than 90% of the column or row total

Example 6 – Group disclosure



Whether ever incarcerated, by selected occupations

	Ever incarcerated (No.)		Ever incarcerated (Row %)	
Occupation Code	Yes	Νο	Yes	Νο
Plumber	12	200	6%	94%
Sales Assistant	110	102	52%	48%
Police officer	0	36	0%	100%
Librarian	140	11	93%	7%

Secondary contributor rules



WHY? Designed to protect the confidentiality where data has been collected and output about one unit (primary contributor) but could disclose information about a higher-level unit (secondary contributor)

WHERE? Output from multi-level datasets

At least 5 businesses or 10 households In addition to the Rule of 10 for the primary contributor

Example 7 – Secondary contributors



- Number of persons per SA3 working full time in the mining industry
- Source: Employee, Earnings and Hours Survey

Area	Total Employees (weighted)
North	10,345
South	5,023
East	44,553
West	24,344
Mid	701

Example 7 – Secondary contributors



- Number of persons per SA3 working full time in the mining industry
- Source: Employee, Earnings and Hours Survey

Area	Total Employees (weighted)	Total Persons (unweighted)	Number of unique Businesses
North	10,345	1057	7
South	5,023	543	2
East	44,553	4754	13
West	24,344	2489	12
Mid	701	65	1

Other outputs





- Charts/graphs supply underlying counts
- Indexes Explain index construction
- Code remove counts and other data

Help us to clear to your outputs quickly



- Checking your output meets the rules and applying treatments
- Clearly labelling and formatting your output
- Providing the required **supporting data**
- Copying both outputs and evidence to your O:/Output drive

Help us to clear to your outputs quickly



- Requesting clearance in a **new email** chain through the link on the website
- Providing detailed descriptions in each field of the output template

Do not put counts or other data into emails

Outputs from the DataLab



- We are human, we make mistakes
 - Inform us if we have made a mistake in clearing your output
 - Don't use files that have been cleared incorrectly
 - Delete files and emails when requested
- Mistakes are investigated for potential breaches and if found to be a breach will be treated accordingly.

Questions and support

Use information on the ABS website


There are rules, and examples plus this learning material.

DataLab User Guide

https://www.abs.gov.au/statistics/microdata-tablebuilder/datalab

DataLab enquiries

Go to "contact us" in the user guide and choose the template that matches your query



ABS DATA INTEGRATION & DIGITAL SERVICES

Bureau of

Statistics

What's next myDATA portal



- Login to the myDATA portal and download the quiz and <u>all</u> the forms
 - Complete the quiz, read, sign and submit all these via email:
 - to: <u>data.services@abs.gov.au</u>
 - subject line: DataLab training quiz and forms

Accessing the DataLab from the User Guide





Australian Bureau of Statistics	Statistics Census Participating in a survey About Q
Home > Statistics > Microdata and TableBuild	
Sections Search this release Q DataLab	DataLab Analyse the most detailed microdata in the secure DataLab for your statistical research or modelling, find out about charges and how to access
On this page What is DataLab	Released 4/11/2021
Charges Applying for DataLab access Enabling access to DataLab myDATA Beta User Portal	Log into DataLab
Privacy policy Topics	What is DataLab DataLab is the analysis solution for high-end users who want to undertake real time complex analysis of detailed
Safe researcher training	microdata. <u>Compare data services</u> to see if detailed microdata in the DataLab is the right service for you.

ABS DATA INTEGRATION & DIGITAL SERVICES



ABS DATA INTEGRATION & DIGITAL SERVICES

ABS DATA INTEGRATION & DIGITAL SERVICES

DataLab

DataLab Safe Researcher Training

Thank you for attending today's training